


Error propagation at the microMott

Greg Blume¹

¹Department of Physics, Old Dominion University, Norfolk VA, 23529 USA

The microMott



- Counting experiment to measure an asymmetry between scattering angles
- Retarding field grids isolate elastic scatterings
- Asymmetry used to extract beam polarization

$$A = P_b S(\theta) \Rightarrow P_b = \frac{A}{S(\theta)}$$

The data

- L/R are left and right detector, $+/-$ are for plus and minus helicity (HWP reversal)

	1	L+	L-	R+	R-	V	
Dark, d_1	2	20.000	20.000	15.000	15.000	152.806	
3x Light, l_i	3	759853.000	647536.000	264401.000	375516.000	152.781	
	4	745859.000	626904.000	259332.000	372740.000	152.810	
Dark, d_2	5	709136.000	610520.000	257920.000	370600.000	152.828	
	6	31.000	31.000	19.000	19.000	152.715	
	7	33.000	33.000	16.000	16.000	163.041	
	8	657430.000	563196.000	220662.000	315151.000	163.030	
	9	...					

Retarding Field at 150 V


- Retarding Field scanned from 150 to 320 V to include threshold voltage (248 V)

The data reduction

- Work now in cells of one retarding field voltage and only L+
- Remove dark count average from each I_i

$$I_i^* = I_i^* \pm \Delta d = I_i - \bar{d} \pm \Delta d$$

- Remaining counts above threshold are x-rays, remove them




The data reduction

- Extrapolate to voltages below threshold
- Define the background $I_{bg}^{(i)}$ for each voltage and subtract to produce adjusted spectra

$$I_{bg}^{(i)} \pm \Delta I_{bg}^{(i)} = (mv_i + b) \pm \sqrt{(v_i \Delta m)^2 + (\Delta b)^2}$$

$$c_i \pm \Delta c_i = I_i^* - I_{bg}^{(i)} \pm \Delta I_{bg}^{(i)}$$



The asymmetry

- Calculate asymmetry for each v_i in the cell — our data file now looks like below where $l_{pi} = c_i$ for each column

$$A_i = \frac{1 - \sqrt{r_i}}{1 + \sqrt{r_i}}, \quad r = \frac{N_i^-}{N_i^+}, \quad N_i^- = l_{mi}r_{pi}, \quad N_i^+ = l_{pi}r_{mi}$$

1	L+	L-	R+	R-	V
2	lp1	lm1	rp1	rm1	v1
3	lp2	lm2	rp2	rm2	v2
4	lp3	lm3	rp3	rm3	v3
5	lp1	lm1	rp1	rm1	v1
6	...				

$$\Rightarrow \Delta N_i^- = (l_{mi}r_{pi}) \sqrt{\left(\frac{\Delta l_{mi}}{l_{mi}}\right)^2 + \left(\frac{\Delta r_{pi}}{r_{pi}}\right)^2}$$

$$\Rightarrow \Delta N_i^+ = (l_{pi}r_{mi}) \sqrt{\left(\frac{\Delta l_{pi}}{l_{pi}}\right)^2 + \left(\frac{\Delta r_{mi}}{r_{mi}}\right)^2}$$

$$\Rightarrow \Delta r_i = \frac{N_i^-}{N_i^+} \sqrt{\left(\frac{\Delta N_i^-}{N_i^-}\right)^2 + \left(\frac{\Delta N_i^+}{N_i^+}\right)^2} \Rightarrow \Delta A_i = \frac{A_i \Delta r_i}{\sqrt{2}} \sqrt{\frac{r_i + 1}{r_i(r_i - 1)^2}}$$

The asymmetry


- Each Voltage cell now has the form

1	A	dA	V
2	A1	dA1	v1
3	A2	dA2	v2
4	A3	dA3	v3
5	A1	dA1	v1
6	...		

- Calculate the average Asymmetry (and Voltage) for threshold extrapolation

$$A \pm \Delta A = \bar{A} \pm \frac{\sqrt{\sum_i (\Delta A_i)^2}}{3}, \quad V = \bar{V}$$

- Cells are condensed to one asymmetry per cell, use 6 cells



Weighted fit applied
Error grows statistically
 $y = (s \pm \Delta s)x + (i \pm \Delta i)$

The polarization


- Take value of fit at 248 V and divide by $S(\theta) = 0.201$

$$A_{248} \pm \Delta A_{248} = (s(248) + i) \pm \sqrt{(\Delta s(248))^2 + (\Delta i)^2}$$
$$\implies P \pm \Delta P = \frac{A_{248}}{S(\theta)} \pm \frac{\Delta A_{248}}{S(\theta)}$$


- Example polarization result is **78.04 ± 6.67 %**
- Does not include error on the Sherman function
- How does this stack up against old method? What changed?

The comparison

Old results



New results



- Old polarization result is **78.71 ± 3.37 %**
- Error bars are **not** statistical, were calculated by the standard deviation of the asymmetries
- **No** error propagation from the counting statistics
- Fit was **unweighted** — doesn't account for random errors from σ

The takeaways

- Not accounting for error propagation from the counts discards information — cannot ensure statistical behavior
- Using the standard deviation for the error w/ an unweighted fit can underestimate the error
- Weighted fits are important to capture the statistical behavior of a counting experiment
- Good statistics are **IMPORTANT**, otherwise error is large
- On another run, polarization is 83.6772 ± 19.761 %, and error only gets worse
- **NEED** to have > 200000 events for acceptable statistics