

Darklight Omesie 1c at the CEBAF injector

Jan C. Bernauer

April 11, 2018




Massachusetts Institute of Technology

^8Be is special

Many images from arXiv:1707.09749

^8Be is special: two **narrow, highly energetic states** which can decay to ground state via **E/M**




Decay modes of $^8\text{Be}(18.15)$





Hadronic, electromagnetic and through **internal pair conversion**

The Atomkin experiment



1.04 MeV proton beam on ${}^7\text{Li}$ to ${}^8\text{Be}(18.15) + \gamma$. Followed by decay. Looked at e^\pm pairs from internal conversion.

The beryllium anomaly



Why believe it?

- This model has $\chi^2/d.o.f.$ of 1.07, significance of 6.8σ
- Bump, not last bin effect
- Rises/falls when scanning through proton energies around resonance
- Excess only happens for symmetric-energy pairs
- Preliminary reports of same excess in ${}^8\text{Be}(17.6)$ (same group)

Why not believe it?

- Group has a history of finding peaks
- IIUC, the detector acceptance has a minimum at 140°
- DM boson interpretation is proto-phobic to evade NA48/2 limits
 - Actually: $\frac{\epsilon_p}{\epsilon_n}$ coupling below $\pm 8\%$. Z^0 is $\sim 7\%$

We can measure it!

In DarkLight, production is via **Bremsstrahlung**,
predominantly **ISR off the electron**.

We can look at $e^- p \rightarrow e^- p X$, followed by $X \rightarrow e^- p (e^- e^+)$

Irreducible background: $e^- p \rightarrow e^- p \gamma^* \rightarrow e^- p e^+ e^-$

We can measure it!

In DarkLight, production is via **Bremsstrahlung**, predominantly **ISR off the electron**.

We can look at $e^- p \rightarrow e^- p X$, followed by $X \rightarrow e^- p (e^- e^+)$

Irreducible background: $e^- p \rightarrow e^- p \gamma^* \rightarrow e^- p e^+ e^-$

Best kinematics:

- highest production rate if X takes all electron energy.
CS rise beats all
- with limited out-of-plane acceptance, **symmetric angle optimal**

- Replace hydrogen target with tantalum. Point like, and more luminosity
- **Main background is NOT the irreducible one.** Random coincidences between
 - radiative elastic electrons
 - positrons from (virtual) photon pair-production where e^- is missed
- Can optimize by moving electron arm backward

Proposed setup

- 45 MeV beam, 150 μA on 10 μm tantalum foil
→ about 0.3 inv. fb/s hydrogen equivalent
- Positron spectrometer at 16° , 28 MeV
- Electron spectrometer at 33.5° , 15 MeV

Spectrometer design parameters

Kinematic var.	Acc.	Inv. mass res.	est. res. on focal plane	Error
in-plane angle	$\pm 2^\circ$	$22 \frac{\text{keV}}{\text{mrad}}$	5mm/7cm \rightarrow 1.4 mrad	32 keV
out-of-plane angle	$\pm 5^\circ$	$5 \frac{\text{keV}}{\text{mrad}}$	1° and $1.14^\circ \rightarrow 1.5^\circ$	133 keV
momentum	$\pm 20\%$	$85 \frac{\text{keV}}{\%}$	5mm/30cm \rightarrow $< 0.2\%$	17 keV

Sum for two spec: 194 keV, assumed 250 keV

QED irreducible: 55 Hz coincidences,

QED irreducible: 55 Hz coincidences,
... but 120 kHz e^+ singles

Background rates

QED irreducible: 55 Hz coincidences,

... but 120 kHz e^+ singles


Elastic e^- + internal Bremsstrahlung: 6 MHz

→ Random coincidence rate 500 Hz


(at 1.5 GHz bunch rate)

This is the minimum trigger rate and sets the sensitivity.

Counting rates: A' signal



Counting rates: Backgrounds




Dominated by accidental background


- Random coincidences dominate
- Scaling with instantaneous luminosity:
 - Signal $S \sim \mathcal{L}$
 - QED background $Q \sim \mathcal{L}$
 - Accidental background $A \sim \mathcal{L}^2$
 - Sensitivity $\frac{S}{\sqrt{Q+A}} \sim 1$

Dominated by accidental background

- Random coincidences dominate
- Scaling with instantaneous luminosity:
 - Signal $S \sim \mathcal{L}$
 - QED background $Q \sim \mathcal{L}$
 - Accidental background $A \sim \mathcal{L}^2$
 - Sensitivity $\frac{S}{\sqrt{Q+A}} \sim 1$
- Sensitivity almost independent of luminosity. Scale is set by bunch-clock / time resolution

Space requirements





- Reach plot assumes that we can associate tracks to bunches on the 1.5 GHz level.
Fast trigger detectors!
- Need to study more background sources, but I think we have most.
- 1c can/should be test platform for streaming readout.
What approach works best?