




Bubble Chamber


A novel technique for measuring thermonuclear
rates at low energies

Challenges associated with the measurement of $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$



Count rate estimate
for:

$$\sigma=1 \text{ pb}, I=100 \text{ p}\mu\text{A}, \\ T_{\text{target}}=12 \text{ }\mu\text{g/cm}^2$$




$$\frac{dN}{dt} = \sigma \cdot I \cdot T \cong 1 \text{ count/day}$$

Time Reversal Technique

- $0(1,2)3 \text{ vs } 3(2,1)0 :$
$$\frac{\sigma_{23 \rightarrow 01}}{\sigma_{01 \rightarrow 23}} = \frac{(2j_o + 1)(2j_1 + 1)}{(2j_2 + 1)(2j_3 + 1)} \frac{k_{01}^2}{k_{23}^2}$$

$$^{12}\text{C}(\alpha, \gamma)^{16}\text{O} \text{ vs } ^{16}\text{O}(\gamma, \alpha)^{12}\text{C} : \frac{\sigma_{\gamma, \alpha}}{\sigma_{\alpha, \gamma}} = \frac{2\mu_{\alpha, \gamma}c^2E_{\alpha, \gamma}}{2E_\gamma^2}$$





- The large range of incident γ -rays allows us to use targets with thickness of $\sim 1\text{-}10 \text{ g/cm}^2$

$$L = I \cdot T \cdot \varepsilon \cong 8.4 \times 10^{32}$$

$$R = L \cdot \sigma \cong 70 \text{ counts/day/pb}$$

Bubble Chamber : For Nuclear Astrophysics



Proof of Principle Experiment at H γ S

Case Study :

$^{15}\text{N}(\alpha, \gamma)^{19}\text{F}$ via the time inverse $^{19}\text{F}(\gamma, \alpha)^{15}\text{N}$ process

Astrophysical Motivation :

This reaction is the last link in the thermonuclear reaction chain leading to formation of fluorine in AGB stars

Resonance under study :

$E_x = 5.337 \text{ MeV}, J^\pi = \frac{1}{2}^+$

Target + Buffer Fluid :


$\text{C}_4\text{F}_{10} + \text{H}_2\text{O}$

Superheat conditions :


$T = 30^\circ\text{C}, P = 3 \text{ atm}$

100 Hz Digital Camera

$\Delta t = 10 \text{ ms}$



$$N_\gamma = 2 \times 10^3 - 3 \times 10^6 \text{ } \gamma/\text{sec}$$





Lower Limit of H γ S Measurement

Electron Beam Energy : 400 MeV
Electron Beam Current : 41 mA
Interaction Length: 35m



Strong Bremsstrahlung background
when coupled with large cross-
sections at high energies




Goal of the Experiment at JLab

- First test of the bubble chamber with a Bremsstrahlung beam
- Study the cosmic background level
- Study the background contributions from photodisintegration of nuclei in the superheated N₂O liquid



Background from oxygen isotopes and nitrogen in N₂O

- a) $^{18}\text{O}(\gamma,\alpha)^{14}\text{C}$ (Q-value = -6.23 MeV)
- b) $^{17}\text{O}(\gamma,\alpha)^{13}\text{C}$ (Q-value = -6.36 MeV)
- c) $^{14}\text{N}(\gamma,p)^{13}\text{C}$ (Q-value = -7.55 MeV)
- d) $^{17}\text{O}(\gamma,n)^{16}\text{O}$ (Q-value = -4.14 MeV)




Bubble Distribution


Hl γ S Data




Jlab Data







8 MeV, 0.035 μ A



8 MeV, 0.4 μ A



Results from the Experiment



Conclusions

- ✓ New limit of γ -ray insensitivity of the bubble chamber : 1 in 10^{12} (earlier limit = in 10^9)
- ✓ Cosmic background rate : JLab = 1 in 17 minutes, HLyS = 1 in 2 minutes
- ✓ Rate limit of the bubble chamber is 10^{-3} counts/s at 4 MeV beam energy, we reach this limit at a cross-section of 10 pb for C_4F_{10} target fluid !

Future Plans

- ✓ Study $^{19}F(\gamma,\alpha)^{15}N$ at cross-sections below 3 nb (beam time approved at JLab in August 2016)
- ✓ Study of ^{16}O enrichment : $^{17,18}O < 10^{-6}$

Collaboration



B. DiGiovine
R. J. Holt
K. E. Rehm
R. Talwar



C. Ugalde



A. Robinson
A. Sonnenschein



J. Benesch
J. Grames
G. Kharashvili
D. Meekins
D. Moser
M. Poelker
M. Stutzman
R. Suleiman
C. Tennant




THANK YOU

Back - Up

Bremsstrahlung Beam

$^{16}\text{O}(\gamma, \alpha)^{12}\text{C}$ is an ideal case for a Bremsstrahlung beam :

- Very steep cross-section, only photons near the endpoint contribute to the yield
- No structure (resonances)



Bremsstrahlung spectra calculated using GEANT4 and FLUKA

Penfold-Leiss Cross-section Unfolding

- ❖ Measure yields at electron beam kinetic energy $E = E_1, E_2 \dots, E_n$
- ❖ Yield can be expressed as the convolution of the cross-section with the Bremsstrahlung spectrum :



$$y(E_i) = \int_{threshold}^{E_i} N_\gamma(E_i, k) \sigma(k) dk \approx \sum_{j=1}^i N_\gamma(E_i, \Delta, k_j) \sigma(k_j)$$

Where $N_\gamma(E_i, \Delta, k_j)$ is the number of gammas in the energy bin of width $\Delta = E_i - E_{i-1}$



- ❖ The solution to the above equation gives the cross-section and the corresponding error as follows :

$$\sigma_i = \frac{1}{N_{ii}} [y_i - \sum_{j=1}^{i-1} (N_{ij} \sigma_j)]$$

$$\left(\frac{d\sigma_i}{\sigma_i} \right)^2 = \frac{[(dy_i)^2 + \sum_{j=1}^{i-1} (N_{ij} d\sigma_j)^2]}{[y_i - \sum_{j=1}^{i-1} (N_{ij} \sigma_j)]^2}$$




Role of $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ reaction in Stellar Helium Burning



- It defines the ratio of carbon to oxygen in stellar cores and, as a result, in the universe
- It affects the synthesis of most of the elements of the periodic table
- Determines the minimum mass required by a star to become a supernova


*Suppression of neutron
events by a factor of 100
using acoustic signals*



Traditional Techniques

Assuncao et al., PRC73, 055801 (2006)

9 EUROGAM detectors



Count rate estimate for:

$$\sigma=1 \text{ pb}, I=100 \text{ p}\mu\text{A}, T_{\text{target}}=12 \text{ }\mu\text{g/cm}^2$$



$$\frac{dN}{dt} = \sigma \cdot I \cdot T \cong 1 \text{ count/day}$$

