

Unpolarized positron beam experiment in Hall C

*Proposal to PAC48
based on
LoI to PAC46*

A. Camsonne,¹ M. Carmignotto,¹ R. Ent,¹ J. Grames*,¹ C. Keppel,¹ M. McCaughan,¹
B. Sawatzky,¹ A. Somov,¹ B. Wojtsekhowski,¹ S. Wood,¹ C. Zorn,¹ M. Caudron,²
L. Causse,² P. Chatagnon,² R. Dupré,² M. Ehrhart,² M. Guidal,² S. Habet,² A. Hobart,²
D. Marchand,² C. Muñoz Camacho*†,² S. Niccolai,² H.-S. Ko,² K. Price,² V. Sergeyeva,²
E. Voutier,² S. Zhao,² M. Mazouz*,³ S. Ali,⁴ V. Berdnikov,⁴ T. Horn,⁴ G. Kalicy,⁴
M. Muhoza,⁴ I. Pegg,⁴ R. Trotta,⁴ A. Asaturyan,⁵ A. Mkrtchyan,⁵ H. Mkrtchyan,⁵
V. Tadevosyan,⁵ H. Voskanyan,⁵ S. Zhamkochyan,⁵ M. Amaryan,⁶ C. Hyde,⁶ M. Kerver,⁶
H. Rashad,⁶ J. Murphy,⁷ J. Roche,⁷ P. Markowitz,⁸ A. Afanasev,⁹ W. J. Briscoe,⁹
I. Strakovsky,⁹ M. Boer,¹⁰ R. Paremuzyan,¹⁰ T. Forest,¹¹ J. R.M. Annand,¹² D. J. Hamilton,¹²
B. McKinnon,¹² D. Day,¹³ D. Keller,¹³ R. Rondon,¹³ J. Zhang,¹³ K. Brinkmann,¹⁴ S. Diehl,¹⁴
R. Novotny,¹⁴ P. Gueye,¹⁵ V. Bellini,¹⁶ D. Dutta,¹⁷ E. Kinney,¹⁸ P. Nadel-Turonski,¹⁹
G. Niculescu,²⁰ S. Sirca,²¹ I. Albayrak,²² M. A. I. Fernando,²³ and M. Defurne²⁴

¹Thomas Jefferson National Accelerator Facility
12000 Jefferson Avenue, Newport News, VA 23606, USA

²Laboratoire de Physique des 2 Infinis Irène Joliot-Curie
Université Paris-Saclay, CNRS/IN2P3, IJCLab (Orsay, France)

³Faculté des Sciences de Monastir (Tunisia)

⁴The Catholic University of America
Washington, DC 20064, USA

⁵A. Alikhanyan National Laboratory, Yerevan Physics Institute, Yerevan 375036, Armenia

⁶Old Dominion University
Norfolk, VA 23529, USA

⁷Ohio University
Athens, OH 45701, USA

⁸Florida International University
Miami, FL 33199, USA

⁹The George Washington University
Washington, DC 20052, USA

¹⁰University of New Hampshire
Durham, NH 03824, USA

¹¹Idaho State University
Pocatello, ID 83209, USA

¹²University of Glasgow
Glasgow G12 8QQ, United Kingdom

¹³University of Virginia
Charlottesville, VA 22904, USA

¹⁴Universität Gießen
Ludwigstraße 23, 35390 Gießen, Deutschland

¹⁵Facility for Rare Isotope Beams, Michigan State University
640 South Shaw Lane, East Lansing, MI 48824

¹⁶Istituto Nazionale di Fisica Nucleare

Sezione di Catania, 95123 Catania, Italy

¹⁷Mississippi State University
Mississippi State, MS 39762, USA

¹⁸University of Colorado
Boulder, CO 80309, USA

¹⁹Stony Brook University
Stony Brook, NY

²⁰James Madison University
Harrisonburg, VA 22807, USA

²¹Univerza v Ljubljani
1000 Ljubljana, Slovenia

²²Akdeniz Üniversitesi
07070 Konyaalti/Antalya, Turkey


²³Hampton University
Hampton, VA 23668

²⁴Commissariat à l'Energie Atomique
91191 Gif-sur-Yvette, France


* Spokesperson

† Contact person



Deeply Virtual Compton Scattering (DVCS)




$$ep \rightarrow ep\gamma =$$




Calculable in QED with our ~1% knowledge of form factors at low momentum transfer



3D imaging of the proton with DVCS



Motivation for using positrons



$$|\mathcal{T}(\pm ep \rightarrow \pm ep \gamma)|^2 = |\mathcal{T}^{BH}|^2 + |\mathcal{T}^{DVCS}|^2 \mp \mathcal{I}$$

Opposite sign
for e^- & e^+

When only 1 quark of the proton is involved in the reaction:

$$\begin{aligned} d^5 \vec{\sigma} - d^5 \overleftarrow{\sigma} &= \Im(T^{BH} \cdot T^{DVCS}) \\ d^5 \vec{\sigma} + d^5 \overleftarrow{\sigma} &= |BH|^2 + \Re(T^{BH} \cdot T^{DVCS}) + |DVCS|^2 \end{aligned}$$

DVCS program at JLab

Two complementary approaches:

- Survey measurements with large acceptance device (CLAS + CLAS12):

Study of many different observables over a wide range of kinematics,
but limited statistical and systematic uncertainties

- Precision measurements in selected kinematic settings (Hall A + Hall C):

test of scaling, higher twist corrections, L/T separations...


A few milestones of the precision DVCS program

- First indications of leading twist dominance for DVCS for Q^2 as low as $\sim 2 \text{ GeV}^2$
- Large magnitude of the DVCS² contribution
Phys. Rev. Lett. **97**, 262002 (2006)
Phys. Rev. **C92**, 055202 (2015)
- Necessity to include corrections $O(t/Q^2)$ & $O(M^2/Q^2)$ to the DVCS cross section
- *Initial separation DVCS² & BH-DVCS interference (yet ambiguous)*
Nature Communications **8**, 1408 (2017)
- Flavor separation of CFFs combining proton & neutron DVCS data
- DVCS on coherent deuteron (\rightarrow nuclear GPDs)
Phys. Rev. Lett. **99**, 242501 (2007)
Nature Physics **16**, 191 (2020)
- L/T separation of π^0 electroproduction cross section (\rightarrow transversity GPDs)
- Flavor separation of transversity GPDs using π^0 electroproduction & a LD₂ target
Phys. Rev. **C83** 025201 (2011)
Phys. Rev. Lett. **117**, 262001 (2016)
Phys. Rev. Lett. **118**, 222002 (2017)

E07-007: Rosenbluth-like separation of DVCS

$$\sigma(ep \rightarrow ep\gamma) = \underbrace{|BH|^2}_{\text{Known to } \sim 1\%} + \underbrace{\mathcal{I}(BH \cdot DVCS)}_{\text{Linear combination of GPDs}} + \underbrace{|DVCS|^2}_{\text{Bilinear combination of GPDs}}$$

$$\mathcal{I} \propto 1/y^3 = (k/\nu)^3,$$
$$|\mathcal{T}^{DVCS}|^2 \propto 1/y^2 = (k/\nu)^2$$




φ -dependence provides 5 independent observables:

$$\sim 1, \sim \cos \varphi, \sim \sin \varphi, \sim \cos(2\varphi), \sim \sin(2\varphi)$$

E07-007: Rosenbluth-like separation of DVCS

DVCS² and \mathcal{I} (DVCS·BH) separated in NLO and higher-twist scenarios



- DVCS² & \mathcal{I} significantly different in each scenario
- Sizeable DVCS² contribution in the higher-twist scenario in the helicity-dependent cross section

Nature Commun. 8, 1408 (2017)

DVCS with positrons and NPS (proposal to PAC48)

$$|\mathcal{T}(\pm ep \rightarrow \pm ep \gamma)|^2 = |\mathcal{T}^{BH}|^2 + |\mathcal{T}^{DVCS}|^2 \mp \mathcal{I}$$

Opposite sign
for e^- & e^+

Physics goals and motivation:

- ✓ Precise determination of the absolute photon electro-production cross section
- ✓ Clean, model-independent separation of DVCS² and DVCS-BH interference
- ✓ More stringer constraints on theory by combining e^- & e^+ data


In a nutshell:

- Same experimental configuration as approved experiment E12-13-010
- Expected positron beam momentum spread comparable with current electron beam
- Positron beam size larger than current electron beam (twice bigger at 11 GeV according to current simulation)
- No additional systematic uncertainties expected due to the use of positrons

PR12-20-012: Kinematic settings

Same kinematics settings as approved
E12-13-010 with electrons

77 days, 5 μA of (unpolarized) positrons assumed
Positron data: 25% of statistics of electron data




x_{Bj}	0.2				0.36						0.5			0.6			
$Q^2 \text{ (GeV)}^2$	2.0			3.0	3.0			4.0		5.5	3.4		4.8	5.1			6.0
$k \text{ (GeV)}$	6.6	8.8	11		6.6	8.8	11	8.8	11		8.8	11		6.6	8.8	11	
$k' \text{ (GeV)}$	1.3	3.5	5.7	3.0	2.2	4.4	6.6	2.9	5.1	2.9	5.2	7.4	5.9	2.1	4.3	6.5	5.7
$\theta_{\text{Calo}} \text{ (deg)}$	6.3	9.2	10.6	6.3	11.7	14.7	16.2	10.3	12.4	7.9	20.2	21.7	16.6	13.8	17.8	19.8	17.2
$D_{\text{Calo}} \text{ (m)}$	6	4		6	3			4	3	4	3						
$\sigma_{M_X^2} \text{ (GeV}^2\text{)}$	0.17			0.22	0.13		0.12	0.15		0.19	0.09		0.11	0.09			
$I_{\text{beam}} \text{ (}\mu\text{A)}$	5																
Days	1	1	3	1	2	3	2	3	4	13	4	3	7	7	2	7	14

Positron production and transport

Injector modifications

[based on PEPPo experiment:
Phys. Rev. Lett. 116, 214801 (2016)]



Electrons

Area	$\delta p/p$ [$\times 10^{-3}$]	ϵ_x [nm]	ϵ_y [nm]
Chicane	0.5	4.00	4.00
Arc 1	0.05	0.41	0.41
Arc 2	0.03	0.26	0.23
Arc 3	0.035	0.22	0.21
Arc 4	0.044	0.21	0.24
Arc 5	0.060	0.33	0.25
Arc 6	0.090	0.58	0.31
Arc 7	0.104	0.79	0.44
Arc 8	0.133	1.21	0.57
Arc 9	0.167	2.09	0.64
Arc 10	0.194	2.97	0.95
Hall D	0.18	2.70	1.03

Dominated by
damping in the
LINACS

Dominated by
synchrotron
rad. in Arcs

Positrons

Area	$\delta p/p$ [$\times 10^{-3}$]	ϵ_x [nm]	ϵ_y [nm]
Chicane	10	500	500
Arc 1	1	50	50
Arc 2	0.53	26.8	26.6
Arc 3	0.36	19	18.6
Arc 4	0.27	14.5	13.8
Arc 5	0.22	12	11.2
Arc 6	0.19	10	9.5
Arc 7	0.17	8.9	8.35
Arc 8	0.16	8.36	7.38
Arc 9	0.16	8.4	6.8
MYAAT01	0.18	9.13	6.19

At 11 GeV,
after Arc9,
 e^+ beam size
~twice bigger
than e^- beam

Averaging
 ϵ_x and ϵ_y :

$$\sqrt{7.6/1.4} \sim 2.3$$

TAC comments on positron




- The implementation of a multi-Hall, high current, high polarization positron beam at CEBAF raises multiple and complex challenges, as detailed in the TAC report
- If the PAC finds our physics program compelling, our collaboration is ready to engage with the Lab to investigate its feasibility.

TAC conclusion:

In conclusion, while a positron beam upgrade is a major upgrade which will require substantial accelerator physics development, a detailed cost and implementation plan, and expensive changes to the CEBAF accelerator, a multi-Hall positron beam capability could have great potential for a future JLAB 12-GeV science program.

Neutral Particle Spectrometer (NPS)

- 1080 PbWO_4 crystals
- 0.6 Tm sweeping magnet
- F250ADC sampling electronics
- Large opening angle beam pipe
- SHMS as carriage for rotation




Separation of DVCS² and BH-DVCS interference

Projections based on the KM15 model (Kumericki and Mueller, 2015)

$x_B=0.2$,
 $Q^2=2.0 \text{ GeV}^2$

$x_B=0.3$,
 $Q^2=4.0 \text{ GeV}^2$

$x_B=0.5$,
 $Q^2=3.4 \text{ GeV}^2$




Systematic uncertainties

Source	pt-to-pt (%)	scale (%)
Acceptance	0.4	1.0
Electron PID	<0.1	<0.1
Efficiency	0.5	1.0
Electron tracking	0.1	0.5
Charge	0.5	1.0
Target thickness	0.2	0.5
Kinematics	0.4	<0.1
Exclusivity	1.0	2.0
π^0 subtraction	0.5	1.0
Radiative corrections	1.2	2.0
Total	1.8-1.9	3.4-3.5

The π^0 electroproduction cross section would be measured concurrently with DVCS with both electrons and positrons, and would allow to monitor the systematics of the e- and e+ runs

Impact on Compton Form Factors (CFFs) extraction



✓ Combined fit of all electron data from approved experiment E12-13-010

(helicity-dependent AND helicity-independent cross sections)


✓ Fits with and without the proposed positron data

In order to extract the CFFs we exploit the combined

- Azimuthal dependence (ϕ)
- Beam-energy dependence
- Q^2 -dependence
- Helicity dependence (for E12-13-010 data)
- **Beam-charge dependence**

of the DVCS cross section

Impact on Compton Form Factors (CFFs) extraction




A factor of 4-6 improvement in the extraction of LO/LT CFFs $\Re e(H)$ and $\Re e(\tilde{H})$
(factor of ~ 2 for HT and NLO)

Correlation coefficients

Correlations between different CFFs are significantly improved by a combined fit with positrons


$$|\rho_{i,j}| = |\text{COV}[\mathbb{F}_i, \mathbb{F}_j] / (\sigma_i \sigma_j)|$$

Electrons only



($t = -0.26 \text{ GeV}^2$)

Electrons & Positrons



LT/LO

HT

NLO

Much better separation of
H & Ht CFFs at LT/LO

Summary and conclusion

- **Positrons** are the **unique way to unambiguously separate** the DVCS² and the BH-DVCS interference
- They will have a strong impact on fits of DVCS data, and the **3D-imaging program of the nucleon**
- **77 days** of (unpolarized) positrons at $I \geq 5$ mA were requested to the PAC
- Same setup (HMS+NPS) and kinematics of approved experiment E12-13-010

Back-up

E07-007: Rosenbluth-like separation of DVCS

- Cross section measured at 2 beam energies and constant Q^2 , x_B , t

$E = 4.5 \text{ GeV}$

$E = 5.6 \text{ GeV}$


$Q^2 = 1.75 \text{ GeV}^2$

$x_B = 0.36$

$t = -0.30 \text{ GeV}^2$

Helicity-independent
cross section


Helicity-dependent
cross section



- Leading-twist and LO simultaneous fit of both beam energies (dashed line) does not reproduce the data

E07-007: Rosenbluth-like separation of DVCS

- Cross section measured at 2 beam energies and constant Q^2 , x_B , t



- Using only helicity-conserving CFFs ("LT/LO") the fit of both beam energies (dashed line) does not reproduce the data
- Including helicity-flip CFFs, either single-helicity flip ("HT") or double-helicity flip ("NLO") satisfactorily reproduce the angular dependence (blue solid line)