


DVCS with positrons and NPS (proposal to PAC48)

Physics goals and motivation:

- ✓ Precise determination of the absolute photon electro-production cross section
- ✓ Clean separation of DVCS² and DVCS-BH interference
- ✓ More stringer constraints on CFFs by combining e^- & e^+ data

Same kinematics settings as approved
E12-13-010 with electrons




x_{Bj}	0.2				0.36						0.5			0.6			
Q^2 (GeV) ²	2.0			3.0	3.0			4.0		5.5	3.4		4.8	5.1			6.0
k (GeV)	6.6	8.8	11		6.6	8.8	11	8.8	11		8.8	11		6.6	8.8	11	
k' (GeV)	1.3	3.5	5.7	3.0	2.2	4.4	6.6	2.9	5.1	2.9	5.2	7.4	5.9	2.1	4.3	6.5	5.7
θ_{Calo} (deg)	6.3	9.2	10.6	6.3	11.7	14.7	16.2	10.3	12.4	7.9	20.2	21.7	16.6	13.8	17.8	19.8	17.2
D_{Calo} (m)	6	4		6	3			4	3	4	3						
$\sigma_{M_X^2}$ (GeV ²)	0.17			0.22	0.13		0.12	0.15		0.19	0.09		0.11	0.09			
I_{beam} (μA)	5																
Days	1	1	3	1	2	3	2	3	4	13	4	3	7	7	2	7	14

77 days, $>5 \mu\text{A}$ of positrons (unpolarized)
Positron data: 25% of statistics of electron data

Positron production and transport

Injector modifications
(based on PEPPo experiment)



Electrons

Area	$\delta p/p$ [$\times 10^{-3}$]	ϵ_x [nm]	ϵ_y [nm]
Chicane	0.5	4.00	4.00
Arc 1	0.05	0.41	0.41
Arc 2	0.03	0.26	0.23
Arc 3	0.035	0.22	0.21
Arc 4	0.044	0.21	0.24
Arc 5	0.060	0.33	0.25
Arc 6	0.090	0.58	0.31
Arc 7	0.104	0.79	0.44
Arc 8	0.133	1.21	0.57
Arc 9	0.167	2.09	0.64
Arc 10	0.194	2.97	0.95
Hall D	0.18	2.70	1.03

Dominated by
damping in the
LINACS

Dominated by
synchrotron rad.
in Arcs

Positrons

Area	$\delta p/p$ [$\times 10^{-3}$]	ϵ_x [nm]	ϵ_y [nm]
Chicane	10	500	500
Arc 1	1	50	50
Arc 2	0.53	26.8	26.6
Arc 3	0.36	19	18.6
Arc 4	0.27	14.5	13.8
Arc 5	0.22	12	11.2
Arc 6	0.19	10	9.5
Arc 7	0.17	8.9	8.35
Arc 8	0.16	8.36	7.38
Arc 9	0.16	8.4	6.8
MYAAT01	0.18	9.13	6.19

Separation of DVCS² and BH-DVCS interference

Projections based on the KM15 model (Kumericki and Mueller, 2015)


$x_B=0.2$,
 $Q^2=2.0 \text{ GeV}^2$

$x_B=0.3$,
 $Q^2=4.0 \text{ GeV}^2$

$x_B=0.5$,
 $Q^2=3.4 \text{ GeV}^2$


Impact on Compton Form Factors (CFFs) extraction

- ✓ Combined fit of all electron data from approved experiment E12-13-010 (helicity-dependent AND helicity-independent cross sections)
- ✓ Fits include LO & LT CFFs, but also +1 helicity-flip CFFs ("HT") and +2 helicity-flip CFFs ("NLO")
- ✓ Cross sections generated with CFFs values fitted to 6 GeV data



Fits and analysis by M. Mazouz (U. of Monastir)

DVCS with positrons and NPS (proposal to PAC48)




A factor of 4-6 improvement in the extraction of LO/LT CFFs $\Re(H)$ and $\Re(\tilde{H})$
(factor of ~ 2 for HT and NLO)

Correlation coefficients (t=-0.26 GeV²)


Correlations between different CFFs are significantly improved by a combined fit with positrons

$$|\rho_{i,j}| = |\text{COV}[\mathbb{F}_i, \mathbb{F}_j] / (\sigma_i \sigma_j)|$$

Electrons only



Electrons & Positrons



LT/LO

HT

NLO

Much better separation of
H & Ht CFFs at LT/LO

Summary and timeline

- Positrons will help to cleanly separate DVCS² and BH-DVCS interference
- Strong impact on CFFs fits and extraction
- Proposal draft available [here](#)
(an updated version will be circulated next week by email)
- PAC deadline: June 22