
Pion and Kaon Structure on the Lattice

David Richards (Jefferson Lab)

PIEIC 2018, CUA, 24-25 May 2018


Outline

- Pion and Kaon EM Form Factors
 - Experimental Motivation
 - Direct Calculation in Lattice QCD
 - Approach to Partonic Degrees of Freedom
- Pion and Kaon PDFs
 - Why the pion?
 - Good Lattice Cross sections
- Future Opportunities

Pion and Kaon Form Factors

Pion EM form factor

Paradigm for LQCD Calculations of matrix elements




$$\langle \pi(\vec{p}_f) | V_\mu(0) | \pi(\vec{p}_i) \rangle = (p_i + p_f)_\mu F(Q^2)$$

where

$$V_\mu = \frac{2}{3} \bar{u} \gamma_\mu u - \frac{1}{3} \bar{d} \gamma_\mu d$$
$$-Q^2 = [E_\pi(\vec{p}_f) - E_\pi(\vec{p}_i)]^2 - (\vec{p}_f - \vec{p}_i)^2$$

Spacelike...

Pion Experimental Summary



Thanks to
Bipasha Chakraborty

G. Huber, D.Gaskell, T. Horn
PR12-16-003

Charge Radius

Partonic DOF

$Q^2 \longrightarrow 30 \text{ GeV}^2$ at future EIC

Anatomy of Pion Form Factor Calculation


$$\Gamma_{\pi^+\mu\pi^+}(t_f, t; \vec{p}, \vec{q}) = \sum_{\vec{x}, \vec{y}} \langle 0 | \phi(\vec{x}, t_f) V_\mu(\vec{y}, t) \phi^\dagger(\vec{0}, 0) | 0 \rangle e^{-i\vec{p} \cdot \vec{x}} e^{-i\vec{q} \cdot \vec{y}},$$

Resolution of unity – insert states


$$\langle 0 | \phi(0) | \pi, \vec{p} + \vec{q} \rangle \langle \pi, \vec{p} + \vec{q} | V_\mu(0) | \pi, \vec{p} \rangle \langle \pi, \vec{p} | \phi^\dagger | 0 \rangle e^{-E(\vec{p})(t-t_i)} e^{-E(\vec{p}+\vec{q})(t_f-t)}$$

$$\Gamma_{\pi^+\pi^+}(t, 0; \vec{p}) = \sum_{\vec{x}} \langle 0 | \phi(\vec{x}, t_f) \phi^\dagger(0) | 0 \rangle e^{-i\vec{p} \cdot \vec{x}}$$

$$\propto |\langle 0 | \phi(0) | \pi, \vec{p} \rangle|^2 e^{-E(\vec{p})t}$$




Pion Form Factor - I



$$\langle r^2 \rangle = 6 \left. \frac{dF(q^2)}{dq^2} \right|_{q^2=0}$$

Briceno, Chakraborty, Edwards, Kusno, Orginos, DGR, Winter



Pion Form Factor - II

- Challenge to reach high momenta
 - discretization errors $p \leq 1/a$
- Signal-to-noise ratio

$$C(t, \vec{p}) \equiv \sum_{\vec{x}} \langle 0 | \mathcal{O}(t, \vec{x}) \mathcal{O}^\dagger(0, 0) | 0 \rangle e^{-i\vec{p} \cdot \vec{x}} \rightarrow e^{-E(\vec{p})t}$$

$$C_{\sqrt{\sigma^2}}(t, \vec{p}) \xrightarrow{\vec{x}} e^{-m_\pi t}$$

*Probe correlators
at small t*

Boosted interpolating
operators

Bali et al., Phys. Rev. D
93, 094515 (2016)

Variational Method

Feynman-Hellmann method

Variational Method

- Solve generalized eigenvalue equation

$$C(t)v^{(N)}(t, t_0) = \lambda_N(t, t_0)C(t_0)v^{(N)}(t, t_0).$$

$$\lambda_N(t, t_0) \longrightarrow e^{-E_N(t-t_0)},$$

- Find *optimal interpolating operator*, coupling to lowest state

$$\mathcal{O}_{N,\text{proj}} = v_i^{(N)} \mathcal{O}_i$$

- Implement using *distillation* M.Peardon et al., arXiv:0905.2160

$$C_{3\text{pt}} \rightarrow \langle 0 | \mathcal{O}_{\text{proj}}(\vec{p}_f, t_f) V_\mu(\tau) \mathcal{O}_{\text{proj}}(\vec{p}_i, t_i) | 0 \rangle; \vec{q} = \vec{p}_f - \vec{p}_i$$


Feynman-Hellmann method

$$H = H_0 + \lambda H_\lambda$$

$$\frac{\partial E_n}{\partial \lambda} = \langle n | H_\lambda | n \rangle$$

Reduces to calculation of energy-shift of two-point functions **but** repeat the calculation for each operator

Form Factor at high Q^2



arXiv:1702.01513

arXiv:1710.07554

Form factor at high
momenta achievable

Pion and Kaon PDFs

*J. Karpie, C. Egerer, J.W. Qiu, B. Chakraborty, R.
Edwards, K. Orginos, DGR, R. Sufian*

Pion and Kaon PDFs

- Euclidean lattice precludes the calculation of light-cone correlation functions
 - So... Use *Operator-Product-Expansion* to formulate in terms of *Mellin Moments*

$$q(x, \mu) = \int \frac{d\xi^-}{4\pi} e^{-ix\xi^- P^+} \langle P | \bar{\psi}(\xi^-) \gamma^+ e^{-ig \int_0^{\xi^-} d\eta^- A^+(\eta^-)} \psi(0) | P \rangle$$

↓

$$\langle P | \bar{\psi} \gamma_{\mu_1} (\gamma_5) D_{\mu_2} \dots D_{\mu_n} \psi | P \rangle \rightarrow P_{\mu_1} \dots P_{\mu_n} a^{(n)}$$

- Moment Methods
 - Extended operators: Z.Davoudi and M. Savage, PRD 86,054505 (2012)
 - Valence heavy quark: W.Detmold and W.Lin, PRD73, 014501 (2006)

KF Liu, SJ Dong, PRL72, 1790 (1994)

- Hadronic Tensor (HT) $W_{\mu\nu} = \frac{1}{4\pi} \int d^4z e^{iq \cdot z} \langle p | J_\mu(z)^\dagger J_\nu(0) | p \rangle$
- $$C_4(\vec{p}, \vec{q}, \tau) = \sum_{\vec{x}_f} e^{-i\vec{p} \cdot \vec{x}_f} \sum_{\vec{x}_2, \vec{x}_1} e^{-i\vec{q} \cdot (\vec{x}_2 - \vec{x}_1)} \langle N(\vec{x}_f, t_f) J_\mu(\vec{x}_2, t_2) J_\nu(\vec{x}_1, t_1) \bar{N}(\vec{0}, t_0) \rangle$$

This is a **four-point** function.

PDFs - II

- Quasi-PDF (**qPDF**) interpreted in **LaMET** (Large Momentum Effective Theory) was proposed by X.Ji **X. Ji, Phys. Rev. Lett. 110 (2013) 262002**

$$q(x, \mu^2, P^z) = \int \frac{dz}{4\pi} e^{izk^z} \langle P | \bar{\psi}(z) \gamma^z e^{-ig \int_0^z dz' A^z(z')} \psi(0) | P \rangle + \mathcal{O}((\Lambda^2/(P^z)^2), M^2/(P^z)^2)$$

Quasi distributions approach light-cone distributions in limit of large P^z

$$q(x, \mu^2, P^z) = \int_x^1 \frac{dy}{y} Z\left(\frac{x}{y}, \frac{\mu}{P^z}\right) q(y, \mu^2) + \mathcal{O}(\Lambda^2/(P^z)^2, M^2/(P^z)^2)$$

- Pseudo-PDF (**pPDF**) recognizing generalization of PDFs in terms of *loffe* Time. $\nu = p \cdot z$

A. Radyushkin, PLB767 (2017)

$$\mathcal{M}^\alpha(z, p) = \langle p | \bar{\psi}(z) \gamma^\alpha \exp\left(-ig \int_0^z dz' A^z(z')\right) \psi(0) | p \rangle$$

Lattice Cross Sections

- Good “Lattice Cross Sections” (**LCS**) Ma and Qiu, Phys. Rev. Lett. 120 022003

$$\sigma_n(\omega, \xi^2, P^2) = \langle P | T\{\mathcal{O}_n(\xi)\} | P \rangle \quad \text{Expressed in coordinate space}$$

where

$$\sigma_n(\omega, \xi^2, P^2) = \sum_a \int_{-1}^1 \frac{dx}{x} f_a(x, \mu^2) K_n^a(x\omega, \xi^2, x^2 P^2, \mu^2) + \mathcal{O}(\xi^2 \Lambda_{\text{QCD}}^2)$$

Calculated in LQCD

Structure function

Calculated in perturbation theory

Short distance scale

Factorize in $\omega = P \cdot \xi, \xi^2 P^2$ providing $\xi \ll \frac{1}{\Lambda_{\text{QCD}}}$

Momentum space

$$\tilde{\sigma}(\tilde{\omega}, q^2 P^2) \equiv \int \frac{d^4 \xi}{\xi^4} \sigma(P \cdot \xi, \xi^2, P^2)$$

$$\tilde{\omega} = 1/x_B$$

Lattice Cross Sections - II

- Quasi- and Pseudo-distributions particular case

$$\mathcal{O}(\xi) = \bar{\psi}(0) \Gamma W(0, 0 + \xi) \psi(\xi)$$

Wilson Line

- Current-current correlators, e.g.


$$\mathcal{O}_S(\xi) = \xi^4 Z_S^2 [\bar{\psi}_q \psi_q](\xi) [\bar{\psi}_q \psi](0)$$

$$\mathcal{O}_{V'}(\xi) = \xi^2 Z_{V'}^2 [\bar{\psi}_q \xi \cdot \gamma \psi_{q'}](\xi) [\bar{\psi}_{q'} \xi \cdot \gamma \psi](0) \\ F_{\mu\rho} F_{\nu}^{\rho}$$

- Gauge-invariant
- Renormalization straightforward

Pion PDFs

- u distribution of FNAL E615 to leading order
- C12-15-006 at Hall A will look at structure of pion
- C12-15-006A at Hall A will look at structure of Kaon
- No free pion target



de Teramond, liu, Sufian, Dosch,
Brodsky, Deur, PRL (2018)

Discrepancy in large- x
behavior of pion distribution

Pion PDFs - II

- Pion less computationally demanding than nucleon
 - *Larger signal-to-noise ratio*



$$C(t, \vec{p}) \equiv \sum_{\vec{x}} \langle 0 | \mathcal{O}(t, \vec{x}) \mathcal{O}^\dagger(0, 0) | 0 \rangle e^{-i\vec{p} \cdot \vec{x}} \rightarrow e^{-E(\vec{p})t}$$

$$C_{\sqrt{\sigma^2}}(t, \vec{p}) \rightarrow \begin{cases} e^{-m_\pi t} & \text{Mesons} \\ e^{-(3m_\pi/2)t} & \text{Baryons} \end{cases}$$



- Important constraint on systematic uncertainty is understanding operator renormalization
 - *Operator renormalization “independent” of external states*
- Admits simple computational methodology
 - *Coordinate-space currents computationally demanding in lattice QCD*

Quasi-Distribution of Pion


$$m_\pi \simeq 300 \text{ MeV}$$



LP3, arXiv:1804.01483



Computational Setup



$$\begin{aligned}
 \langle \Pi(-p') | \mathcal{O}_{J_1}(x_0) \mathcal{O}_{J_2}(\xi) | \Pi(-p') \rangle &= \\
 &= \sum_{y,z} e^{i(p' \cdot z - p \cdot y)} \langle \bar{q} \Gamma_{\Pi} q(z, T) \bar{Q} J_2 Q(x_0 + \xi, t) \bar{q} J_1 q(x_0, t) \bar{q} \Gamma_{\Pi} q(y, 0) \rangle \\
 &= \sum_{y,z} e^{i(p' \cdot z - p \cdot y)} \text{Tr} [J_2 G_Q(x_0 + \xi, t; x_0, t) J_1 G(x_0, t; y, 0) \Gamma_{\Pi} G(y, 0; z, T) \Gamma_{\Pi} G(z, T; x_0 + \xi, t)]
 \end{aligned}$$

Straightforward computational setup using sequential-source method:


$$\begin{aligned}
 D(Z, T; w) H(w; x_0, t) &= \sum_y e^{-ip \cdot y} \Gamma_{\Pi} G(y, 0; x_0, t) \\
 D(s; w) \tilde{H}(w; x_0, t) &= \sum_z e^{ip \cdot z} \Gamma_{\Pi} H(z, T; x_0, t)
 \end{aligned}$$

Preliminary Results

- 2+1 Flavor clover-fermion action

$$a \simeq 0.12, 0.09 \text{ fm}$$

$$m_\pi \simeq 400, 440 \text{ MeV}$$




$$J_1 = V, J_2 = A$$

- 110 configurations
- Single source point for current J_1


Clear isolation of pion matrix element

Preliminary Results - II




Clear signal in real part to p around 1 GeV

Imaginary part only to small values of p



Challenges/Questions

High spatial momentum and lattice systematics



Boosted interpolating operators

Bali et al., Phys. Rev. D 93, 094515 (2016)

Inverse Problem - common to all approaches


$$\sigma_n(\omega, \xi^2, P^2) = \sum_a \int_{-1}^1 \frac{dx}{x} f_a(x, \mu^2) K_n^a(x\omega, \xi^2, x^2 P^2, \mu^2) + \mathcal{O}(\xi^2 \Lambda_{\text{QCD}}^2)$$

Calculate on Lattice


Extract PDF?

Calculate in PQCD

Finite Volume Effects



Briceno, Guerrero, Hansen and Monahan, arXiv:1805.01304



Typically $m_\pi L \simeq 4$



Future? $\left\{ \begin{array}{l} \xi \text{ shortdistance} \\ m_\pi \rightarrow m_\pi^{\text{phys}} \end{array} \right.$

Pion Quark Distribution Amplitude

“Pseudo” quark distribution amplitudes





Bali et al., arXiv:1709.04325

$$T(\omega, z^2) = \langle \pi(\vec{p} \mid [\bar{u}Q](z/2)[\bar{Q}u](-z/2) \mid 0 \rangle$$



Quark Distribution Pictures \longleftrightarrow Ioffe pseudo-time

Pion QDA - II



Fixed z^2

Summary

- Pion Form Factors at high momenta with reach comparable to 12 GeV at JLab
- Calculation of current-current correlators for pion *and kaon* in progress for variety of local operators
 - Important to understand finite-volume effects
 - Extending calculation to close-to-physical

$$m_\pi \simeq 170 \text{ MeV}$$

$$64^3 \times 128 \text{ Lattices}$$

- Variety of lattice cross sections - including pseudo PDFs - on same ensemble of lattices.

Outlook

