

Physics Perspectives for Future K -Long Facility

Igor Strakovsky*


The George Washington University
(for GlueX Collaboration)



- Thermodynamics at freeze-out.
- Spectroscopy of hyperons.
- $K_L p$ data.
- Opportunity with K_L beam.
- Neutron background.
- Expected $K_L p$ data.
- Summary.



History of the Universe




- There is Influence of possible “missing” hyperons on QCD thermodynamics, on **freeze-out** In heavy ion & hadon collisions & in **early Universe**, & in spectroscopy.
- Advance our understanding of **formation of baryons** from **quarks & gluons** **microseconds** after **Big Bang** & in today’s experiments, & connection these developments to experimental searches for direct, spectroscopic, evidence for these “missing” resonances.



Thermodynamics at Freeze-Out

- Recent studies that compare **LQCD** calculations of thermodynamic, statistical **Hadron Resonance Gas** models, & ratios between measured yields of different hadron species in heavy ion collisions provide indirect evidence for the presence of “missing” resonances in all of these contexts.



A. Bazavov *et al* Phys Rev Lett **113**, 072001 (2014)

- Partial **pressure** of strange **mesons**

$$M_1^S = \chi_2^S - \chi_{22}^{BS},$$

$$M_2^S = \frac{1}{12}(\chi_4^S + 11\chi_2^S) + \frac{1}{2}(\chi_{11}^{BS} + \chi_{13}^{BS}),$$

- Partial **pressure** of strange **baryons**

$$B_1^S = -\frac{1}{6}(11\chi_{11}^{BS} + 6\chi_{22}^{BS} + \chi_{13}^{BS}),$$

$$B_2^S = \frac{1}{12}(\chi_4^S - \chi_2^S) - \frac{1}{3}(4\chi_{11}^{BS} - \chi_{13}^{BS}).$$

• + “Missing” Resonances (QM calculations).

• Contribution from **observed Resonances**. 


- Three independent **ratios** start to coincide in crossover region giving identical results only below **chiral crossover temperature** at physical values of quark masses $T_c = 154 \pm 9$ MeV.

Baryon Sector at PDG16



GW *GW Contribution*

C. Patrignani *et al* Chin Phys C **40**, 090001 (2016)



p	$1/2^+$	****	$\Delta(1232)$	$3/2^+$	****	Σ^+	$1/2^+$	****	Ξ^0	$1/2^+$	****	Λ_c^+	$1/2^+$	****
n	$1/2^+$	****	$\Delta(1600)$	$3/2^+$	***	Σ^0	$1/2^+$	****	Ξ^-	$1/2^+$	****	$\Lambda_c(2595)^+$	$1/2^-$	***
$N(1440)$	$1/2^+$	****	$\Delta(1620)$	$1/2^-$	****	Σ^-	$1/2^+$	****	$\Xi(1530)^0$	$3/2^+$	****	$\Lambda_c(2625)^+$	$3/2^-$	***
$N(1520)$	$3/2^-$	****	$\Delta(1700)$	$3/2^-$	****	$\Sigma(1385)$	$3/2^+$	****	$\Xi(1620)^0$	*		$\Lambda_c(2765)^+$	*	
$N(1535)$	$1/2^-$	****	$\Delta(1750)$	$1/2^+$	*	$\Sigma(1480)$	*		$\Xi(1690)^0$	***		$\Lambda_c(2880)^+$	$5/2^+$	***
$N(1650)$	$1/2^-$	****	$\Delta(1900)$	$1/2^-$	**	$\Sigma(1560)$	**		$\Xi(1820)^0$	$3/2^-$	***	$\Lambda_c(2940)^+$	*	
$N(1675)$	$5/2^-$	****	$\Delta(1905)$	$5/2^+$	****	$\Sigma(1580)$	$3/2^-$	**	$\Xi(1950)^0$	***		$\Sigma_c(2455)$	$1/2^+$	****
$N(1680)$	$5/2^+$	****	$\Delta(1910)$	$1/2^+$	****	$\Sigma(1620)$	$1/2^-$	**	$\Xi(2030)^0$	$\geq 3/2^+$	***	$\Sigma_c(2520)$	$3/2^+$	****
$N(1685)$	*		$\Delta(1920)$	$3/2^+$	**	$\Sigma(1660)$	$1/2^+$	***	$\Xi(2250)^0$	**		$\Sigma_c(2800)$	***	
$N(1700)$	$3/2^-$	***	$\Delta(1930)$	$5/2^-$	***	$\Sigma(1670)$	$3/2^-$	****	$\Xi(2370)^0$	**		Ξ_c^+	$1/2^+$	***
$N(1710)$	$1/2^+$	**	$\Delta(1940)$	$3/2^-$	**	$\Sigma(1690)$	$1/2^-$	**	$\Xi(2500)^0$	**		Ξ_c^0	$1/2^+$	***
$N(1720)$	$3/2^+$	**	$\Delta(1950)$	$1/2^+$	**	$\Sigma(1750)$	$1/2^-$	**	$\Xi(2645)^0$	**		Ξ_c^-	$1/2^+$	***
$N(1800)$	$5/2^+$	**	$\Delta(2000)$	$5/2^+$	**	$\Sigma(1770)$	$1/2^+$	**	$\Xi(2770)^0$	**		Ξ_c^0	$1/2^+$	***
$N(1810)$	$1/2^-$	**	$\Delta(2150)$	$1/2^-$	**	$\Sigma(1775)$	$1/2^-$	****	Ω^-	$3/2^+$	*	$\Xi_c(2645)$	$3/2^+$	***
$N(1830)$	$1/2^+$	**	$\Delta(2200)$	$1/2^-$	**	$\Sigma(1840)$	$3/2^+$	**	$\Omega(1370)^-$	*		$\Xi_c(2790)$	$1/2^-$	***
$N(185)$	$7/2^-$	**	$\Delta(2300)$	$9/2^+$	**	$\Sigma(1920)$	$1/2^+$	**	$\Omega(1370)^-$	*		$\Xi_c(2815)$	$3/2^-$	***
$N(1900)$	$3/2^-$	***	$\Delta(2350)$	$5/2^-$	*	$\Sigma(1925)$	$3/2^-$	****	$\Omega(1370)^-$	*		$\Xi_c(2930)$	*	
$N(1990)$	$7/2^-$	**	$\Delta(2390)$	$7/2^+$	*	$\Sigma(1940)$	$3/2^-$	****	$\Omega(1370)^-$	*		$\Xi_c(2980)$	***	
$N(2000)$	$5/2^+$	**	$\Delta(2400)$	$9/2^-$	**	$\Sigma(2000)$	$1/2^-$	*	$\Omega(1370)^-$	*		$\Xi_c(3055)$	**	
$N(2040)$	$3/2^+$	*	$\Delta(2420)$	$11/2^+$	****	$\Sigma(2030)$	$7/2^+$	****	$\Omega(1370)^-$	*		$\Xi_c(3080)$	***	
$N(2060)$	$5/2^-$	**	$\Delta(2750)$	$13/2^-$	**	$\Sigma(2070)$	$5/2^+$	*	$\Omega(1370)^-$	*		$\Xi_c(3123)$	*	
$N(2100)$	$1/2^+$	*	$\Delta(2950)$	$15/2^+$	**	$\Sigma(2080)$	$3/2^+$	**	$\Omega(1370)^-$	*		$\Xi_c(3123)$	*	
$N(2120)$	$3/2^-$	**				$\Sigma(2100)$	$7/2^-$	*	$\Omega(1370)^-$	*		Ω_c^0	$1/2^+$	***
$N(2190)$	$7/2^-$	****	Λ	$1/2^+$	****	$\Sigma(2250)$	***		Ω_c^0	$3/2^+$	***	$\Omega_c(2770)^0$	$3/2^+$	***
$N(2220)$	$9/2^+$	****	$\Lambda(1405)$	$1/2^-$	****	$\Sigma(2455)$	**		Ξ_{cc}^+	*				
$N(2250)$	$9/2^-$	****	$\Lambda(1520)$	$3/2^-$	****	$\Sigma(2620)$	**		Λ_b^0	$1/2^+$	***			
$N(2600)$	$11/2^-$	***	$\Lambda(1600)$	$1/2^+$	***	$\Sigma(3000)$	*		Σ_b^+	$1/2^+$	***			
$N(2700)$	$13/2^+$	**	$\Lambda(1670)$	$1/2^-$	****	$\Sigma(3170)$	*		Σ_b^0	$1/2^+$	***			
			$\Lambda(1690)$	$3/2^-$	**				Ξ_b^0	$3/2^+$	***			
			$\Lambda(1800)$	$1/2^-$	**				Ξ_b^-	$3/2^+$	***			
			$\Lambda(1810)$	$1/2^+$	**				Ξ_b^0	$1/2^+$	***			
			$\Lambda(1820)$	$1/2^+$	**				Ξ_b^-	$1/2^+$	***			
			$\Lambda(1830)$	$5/2^-$	**				Ω_b^-	$1/2^+$	***			
			$\Lambda(1890)$	$3/2^+$	**									
			$\Lambda(2000)$											
			$\Lambda(2020)$	$7/2^+$	*									
			$\Lambda(2100)$	$7/2^-$	****									
			$\Lambda(2110)$	$5/2^+$	***									
			$\Lambda(2325)$	$3/2^-$	*									
			$\Lambda(2350)$	$9/2^+$	***									
			$\Lambda(2585)$	**										

• Pole position in complex energy plane for hyperons has began to be studied only recently (2010), first of all for $\Lambda(1520)$.

- PDG16 has 109 Baryon Resonances (58 of them are 4* & 3*).
- In case of SU(6) x O(3), it would be required 434 baryons, if all revealed multiplets were completed (three 70 & four 56).

Y. Qung *et al* Phys Lett B **694**, 123 (2010)




Baryon Resonances

- **Three light quarks** can be arranged in **6** baryonic **families**, N^* , Δ^* , Λ^* , Σ^* , Ξ^* , & Ω^* .
- **Number of members in family** that can exist is **not arbitrary**.
- If $SU(3)_F$ symmetry of **QCD** is controlling, then:

Octet: N^* , Λ^* , Σ^* , Ξ^*
Decuplet: Δ^* , Σ^* , Ξ^* , & Ω^*



- Number of experimentally identified resonances of each baryon **family** in  summary tables is $16 N^*$, $10 \Delta^*$, $14 \Lambda^*$, $10 \Sigma^*$, $6 \Xi^*$, & $2 \Omega^*$.
- **Constituent Quark** models, for instance, predict existence of no less than $64 N^*$, $22 \Delta^*$ states with **mass** < 3 GeV.

- Seriousness of “**missing-states**” problem is obvious from these numbers.



- To complete $SU(3)_F$ multiplets, one needs no less than $17 \Lambda^*$, $43 \Sigma^*$, $42 \Xi^*$, & $24 \Omega^*$.

B.M.K. Nefkens, πN Newsletter, **14**, 150 (1997)




Very Strange Resonances & Problem of “Missing” States



R. Koniuk and N. Isgur, Phys Rev Lett **44**, 845 (1980)


• Experimental knowledge of hadron spectrum is **incomplete**: more excited states are expected to exist.

• Ω baryon spectrum in **QM**.



U. Löring *et al* Eur Phys J A **10**, 447 (2001)

• Ω baryon spectrum in **LQCD**.



R.G. Edwards *et al* Phys Rev D **87**, 054506 (2013)

Thick frame: **Hybrid states**




W = 1.45 – 5.05 GeV

SAID: <http://gwdac.phys.gwu.edu/>



- Limited number of K_L induced measurements (1961 – 1982)
 2426 $d\sigma/d\Omega$, 348 σ^{tot} , & 115 **P** observables do not allow today to **feel comfortable** with **Hyperon Spectroscopy** results.



- Most of **data** were obtained from **old** low statistics **measurements** with hydrogen

Bubble Chambers.

- Overall systematics** of previous experiments varies between **15% & 35%**.

Energy binning is much broader than hyperon widths.





- There were no measurements using **polarized target**.

It means that there are no **double polarized** observables which are critical for **complete experiment** program.

- We are not aware of any data on **neutron** target.



Data for $K_L p \rightarrow K_S p$




• No $d\sigma/d\Omega$ data are available for $K_L p \rightarrow K_L p$ below $W = 3$ GeV.

• PWA (KSU&GW) predictions at lower & higher energies tend to agree worse with data than in non-strange case.

Courtesy of Mark Manley, KL2016



Data for $K_L p \rightarrow \pi^+ \Lambda$ & $K^- p \rightarrow \pi^0 \Lambda$




• $K^- p \rightarrow \pi^0 \Lambda$ & $K_L p \rightarrow \pi^+ \Lambda$ amplitudes imply that their observables measured at same energy should be identical except for small differences due to isospin-violating mass differences in hadrons.

• Polarized measurements are **tolerable** for **any PWA** solutions.



Data for $K_L p \rightarrow \pi^+ \Sigma^0$ & $K^- p \rightarrow \pi \Sigma$




Courtesy of Mark Manley, KL2016

- Reactions $K_L p \rightarrow \pi^+ \Sigma^0$ & $K_L p \rightarrow \pi^0 \Sigma^+$ are Isospin selective (only $I = 1$ amplitudes are involved) & $K^- p \rightarrow \pi^0 \Sigma^0$ is isospin selective for $I = 0$ whereas reactions $K^- p \rightarrow \pi^- \Sigma^+$ & $K^- p \rightarrow \pi^+ \Sigma^-$ involve both $I = 0$ & $I = 1$ amplitudes.

- New measurements with K_L -beam would lead to better understanding of Σ^* states & help constrain amplitudes for $K^- p \rightarrow \pi \Sigma$ reactions.

- Quality of $K_L p$ data is comparable to that for $K^- p$ data. It would be advantageous to combine $K_L p$ data in a new coupled-channel PWA with available $K^- p$ measurements.

-  lists only **two** results on
BR to $K\Sigma$ $\Lambda(2100)7/2^-$ (BR < 3%)
 $\Sigma(2030)7/2^+$ (BR < 2%).



How to Search for Missing Hyperons


- ▶ New data for inelastic $K_L^0 p$ scattering would significantly improve our knowledge of Σ^* Resonances
- ▶ Very few polarization data are available for any $K_L^0 p$ reactions but are needed to help remove ambiguities in PWAs
- ▶ To search for missing hyperon resonances, we need measurements of production reactions:
 - Σ^* : $K_L^0 p \rightarrow \pi \Sigma^* \rightarrow \pi \pi \Lambda$
 - Λ^* : $K_L^0 p \rightarrow \pi \Lambda^* \rightarrow \pi \pi \Sigma$
 - Ξ^* : $K_L^0 p \rightarrow K \Xi^*, \pi K \Xi^*$
 - Ω^* : $K_L^0 p \rightarrow K^+ K^+ \Omega^*$
- ▶ If such measurements can be performed with good energy & angle coverage & good statistics, then it is very likely that measurements with K_L^0 beams would find several missing hyperon resonances.

Courtesy of Mark Manley, YSTAR2016




Sample of Hunting for Bumps

γK^+ Outlook at GlueX for $\Lambda(1405)$ Line- Shape Measurement



- Measurement may be feasible



$$K_L^0 p \rightarrow \Lambda(1405) \pi^+ \rightarrow \Sigma^{+0} \pi^{-0+} \pi^+$$



Courtesy of Reinhard Schumacher, KL2016



A bit of History

PHYSICAL REVIEW

VOLUME 138, NUMBER 5B

7 JUNE 1965

First paper on subject

Photoproduction of Neutral K Mesons*

S. D. DRELL AND M. JACOB†

Stanford Linear Accelerator Center, Stanford University, Stanford, California

(Received 6 January 1965)

CP-violation (1964)
Hot topic!



Photoproduction of a neutral K -meson beam at high energies from hydrogen is computed in terms of a K^* vector-meson exchange mechanism corrected for final-state interactions. The results are very encouraging for the intensity of high-energy K_2 beams at high-energy electron accelerators. A typical magnitude is $20 \mu\text{b}/\text{sr}$ for a lower limit of the K^0 photoproduction differential cross section, at a laboratory peak angle of 2° , for 15-BeV incident photons.




FIG. 1. K^* exchange in photoproduction.

[Not dominant]

Our motivation in carrying out this calculation is to emphasize the strong suggestion that an intense “healthy” K_2 beam will emerge from high-energy electron accelerators (SLAC in particular) and will be available for detailed experimental studies.

50 $\mu\text{b}/\text{sr}$




FIG. 3. Center-of-mass differential cross section at 10 BeV. Curve (1) gives the Born approximation. Curve (2) is obtained after subtraction of the $j = \frac{1}{2}$ partial wave. Curves (3) and (4) respectively obtained after the $j = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \frac{7}{2}$, and all partial waves have been corrected for absorption in final state. The result is shown as directly obtained from and drawn by the computer.

Courtesy of Mike Albrow, KL2016



A bit of History

The possibility that useful K_L beam could be made at electron synchrotron by photoproduction was being considered, & 1965 prediction for SLAC by Drell & Jacob was optimistic.


8.B.5 Nuclear Physics B23 (1970) 509-524. North-Holland Publishing Company
8.B.6

PHOTOPRODUCTION OF K^0 MESONS FROM PROTONS AND FROM COMPLEX NUCLEI

M. G. ALBROW[†], D. ASTON, D. P. BARBER, L. BIRD^{††},
R. J. ELLISON, C. HALLIWELL, A. E. HARCKHAM^{†††},
F. K. LOEBINGER, P. G. MURPHY, J. WALTERS^{††} and A. J. WYNROE
*Schuster Laboratories, The University of Manchester,
Manchester M13 9PL*

R. F. TEMPLEMAN
*Daresbury Nuclear Physics Laboratory, Daresbury,
Near Warrington, Lancs.*

Received 16 July 1970




"We were at Manchester Univ. close to Daresbury 5 GeV e-synchrotron."

CP-violation

PHYSICAL REVIEW D VOLUME 7, NUMBER 3 1 FEBRUARY 1973

Production of K_L^0 Mesons and Neutrons from Electrons on Beryllium Above 10 GeV*

G. W. Brandenburg, A. D. Brody, W. B. Johnson, D. W. G. S. Leith, J. S. Loos,[†] G. J. Luste,[‡]
J. A. J. Matthews, K. Moriyasu,[§] B. C. Shen,^{||} W. M. Smart,^{**} F. C. Winkelmann, and R. J. Yamartino
Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305
(Received 14 August 1972)



Systematics of particle anti-particle processes through intrinsic property of K-longs



A Letter of Intent to Jefferson Lab PAC-43.



Physics Opportunities with a Secondary K_L^0 Beam at JLab.

Moskov J. Amaryan (spokesperson),^{1,*} Yakov I. Azimov,² William J. Briscoe,³ Eugene Chudakov,⁴ Pavel Degtyarenko,⁴ Gail Dodge,¹ Michael Döring,³ Helmut Haberzettl,³ Charles E. Hyde,¹ Benjamin C. Jackson,⁵ Christopher D. Keith,⁴ Ilya Larin,¹ Dave J. Mack,⁴ D. Mark Manley,⁶ Kanzo Nakayama,⁵ Yongseok Oh,⁷ Emilie Passemar,⁸ Diane Schott,³ Alexander Somov,⁴ Igor Strakovsky,³ and Ronald Workman³

¹Old Dominion University, Norfolk, VA 23529

²Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg 188300, Russia

³The George Washington University, Washington, DC 20052

⁴Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606

⁵The University of Georgia, Athens, GA 30602

⁶Kent State University, Kent, OH 44242

⁷Kyungpook National University, Daegu 702-701, Korea


⁸Indiana University, Bloomington, IN 47405

(Dated: May 14, 2015)

Hyperon Spectroscopy

- We plan to submit a full **Proposal** for **JLab PAC45** in **2017**.

CEBAF Upgrade to 12 GeV




Upgrade Goals

- Accelerator: 6 GeV \Rightarrow 12 GeV
- Halls A,B,C: $e^- < 11$ GeV, $< 100 \mu\text{A}$
- Hall D: e^- 12 GeV $\Rightarrow \gamma$ -beam


Upgrade Status

- Reached 12 GeV in Dec 2015
- Halls A,D: finished
- Halls B,C: about a year to go



Courtesy of Eugene Chudakov, KL2016


Hall D Beam Line Set up for K-longs



$I_e = 5 \mu\text{A}$
 W-radiator = 0.1 R.L.
 Be-target = 1.7 R.L.

- **Electrons** are hitting **W**-radiator.
- **Photons** are hitting **Be**-target.
- **K_Ls** are hitting the **LH₂**-target within **GLueX** setting.

Hall D / GlueX



GLUEX
collisions
experiment

K-long
beam



1/3/2017

HIPS2017, Washington, DC, February 2017

Igor Strakovsky 18



Hall D/GlueX Spectrometer and DAQ


GLUEX

Resolutions

$$h^\pm: \sigma_p/p \sim 1 - 3\%$$

$$\gamma: \sigma_E/E \sim 6\%/\sqrt{E} \oplus 2\%$$

$$\text{Acceptance } 1^\circ < \theta < 120^\circ$$



Detectors

- ▶ CDC, FDC
- ▶ BCAL, FCAL
- ▶ TOF, ST

Plans to add

- ▶ 2017 L3
- ▶ 2018 DIRC



Photoproduction γp 1.5 kHz for a 10 MHz beam; Trigger $\sum E_{CAL} > X$
 GlueX-I 10 MHz/peak: trigger 20 kHz \Rightarrow DAQ \Rightarrow tape 30 kHz spring 2016
 GlueX-II 50 MHz/peak: trigger 100 kHz \Rightarrow DAQ \Rightarrow L3 farm \sim 20 kHz \Rightarrow tape

Courtesy of Eugene Chudakov, YSTAR2016




Expected Neutron Background

- Most important & unpleasant background for K_L comes from **neutrons**.



99% of neutrons associated with $T < 90$ MeV while 0.6% of them are for $T > 125$ MeV.



For **neutron** calculations, we use **MCNP6** transport code.

K-long & Neutron Rate on GlueX LH₂-target



MC @ 12 GeV



Data @ 16 GeV






FIG. 2. Comparison of the neutron and K_2^0 fluxes at the hydrogen bubble chamber for 2° production with 16-GeV electrons. A.D. Brody *et al* Phys Rev Lett **22**, 966 (1969)



← Flux ratio n/K_L

• Delivered with **60 ns** bunch spacing avoids overlap in range of $p = 0.35 - 10.0$ GeV/c.

• With **proton** beam, ratio $n/K_L = 10^3 - 10^4$.




Expected Energy-Resolution


- Mean lifetime of K^- is **12.38** ns ($\tau = 3.7$ m) whereas mean lifetime of K_L is **51.16** ns ($\tau = 15.3$ m).

Thus, it is possible to perform measurements of $K_L p$ scattering at **lower energies** than $K^- p$ scattering due to higher beam flux.

- Momentum measured with **TOF** between **SC** (surrounded LH_2) & **RF** from **CEBAF**.




- Momentum resolution $\Delta p/p$ is growing with momentum: for 1 GeV/c is **1.7%**, for 2 GeV/c is **6%**.



- For $W < 2$ GeV, $\Delta W < 20$ MeV which is suitable to study **Hyperons** with $\Gamma = 30 - 50$ MeV.






Expected Particle Identification


- dE/dx for pK_S .

- Time difference at primary "vertex" for proton hypothesis for pK_S using TOF.

dEdx vs p/M for proton candidates




$t_{\text{tof}} - t_{\text{flight}} - t_{\text{vertex}}$ vs p for proton



dE/dx vs. p/M for K^+ candidates

- dE/dx for $K^+ \pi^0$.



Courtesy of Simon Taylor, KL2016




Expected Cross Sections vs Bubble Chamber Data

• **GlueX** measurements will span $\cos\theta$ from -0.95 to 0.95 in c.m. above $W = 1490$ MeV.

Courtesy of
Simon Taylor, KL2016
Mark Manley, KL2016


- K_L rate is 10^5 K_L /s.
- Uncertainties (statistics only) correspond to **100** days of running time for:

$K_L p \rightarrow K_S p$





Expected
GlueX Data

$K_L p \rightarrow \pi^+ \Lambda$




BC Data





• Prove motivation of **JLab** Proposal *JLab E-03-105*
Pion Photoproduction from Polarized Target for **FROST** Project.

Transverse Polarization **Longitudinal Polarization**



Average ratio of uncertainties of amplitudes
w/o expected **FROST** data.

- Greatest effect naturally requires measurement of all possible quantities as accomplished by **FROST**.

ηp E: I. Senderovich *et al* Phys Lett B **755**, 64 (2016)
 $\pi^+ n$ E: S. Strauch *et al* Phys Lett B **750**, 53 (2015)
More results are coming...

- **KSU&GW** is doing **PWA** including available $K_L p$ & $K^- p$ data plus expected **GlueX** data to show potential impact of new **Hall D** measurements.



Summary

- Here we reviewed what can be learned by studying $K_L p$ scattering leading to **two-body** final states (1st stage).

At later stages, we plan to do $K_L n$ on LD_2 &

$K_L N$ on aka **FROST** with hydrogen & deuterium.

- **JLab K-long Facility** would advance **Hyperon Spectroscopy** & study of **strangeness** in nuclear & hadronic physics.

It may extract very many missing strange states.

To complete $SU(3)_F$ multiplets, one needs no less than

17 Λ^* , 43 Σ^* , 42 Ξ^* , & 24 Ω^* .

- **Discovering** of missing **low-lying hyperon states** would assist in advance our understanding of **formation** of **baryons** from **quarks** & **gluons** microseconds after **Big Bang**.

• Full Proposal is coming for **PAC45** in **2017**, **WELCOME** to **JOIN US**.



Welcome
NEW MEMBERS



Table 1: Comparison of the K_L production yield. The BNL AGS kaon and neutron yields are taken from RSVP reviews in 2004 and 2005. The Project X yields are for a thick target, fully simulated with LAQGSM/MARS15 into the KOPIO beam solid angle and momentum acceptance [68].

Project	Beam energy (GeV)	Target (λ_T)	$p(K_L)$ (MeV/c)	K_L/s (into $500 \mu\text{sr}$)	n/K_L ($E_n > 10 \text{ MeV}$)
BNL AGS	24	1.1 Pt	300–1200	60×10^6	$\sim 1:1000$
Project X	3	1.0 C	300–1200	450×10^6	$\sim 1:2700$


arXiv:1306.5022, arXiv:1306.5009, arXiv:1306.5024

- First stage of Project X aims for neutrinos.
- Proposed K_L beam can be used to study rare decays & CP-violation.
- It may be impossible to use it for Hyperon Spectroscopy because of momentum range & n/K_L ratio.

J-PARC




Japan Proton Accelerator Research Complex



Two beam lines are under operation


K1.1 & High-p beam lines are under construction



Courtesy of Hiroaki Onishi, KL2016



J-PARC Hadron hall extension



Most aggressive scenario:
we expected to have first beam in 2021/2022

Courtesy of Hiroaki Onishi, YSTAR2016






	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026+
--	------	------	------	------	------	------	------	------	------	------	-------

PANDA Phase 0




PANDA Start Setup



PANDA Phase 1

PANDA Hall available




PANDA Full Setup



Installation

PANDA Phase 2



PANDA Phase 3 RESR



Courtesy of James Ritman, YSTAR2016





61 people & 30 talks

Chudakov

Albrow

Richards

Ramos

Zou

Schumacher

Oset

Montgomery

Kamano

Santopinto

Szczepaniak

Mathieu

Passemar

Taylor

Oh

Pennington

Keith

Kohl

Larin

Amaryan

Manley

Filippi

Myhrer

Degtyarenko

Nakayama

Ohnishi

Goity

Mai

Ziegler

Noumi

PHYSICS WITH NEUTRAL KAON BEAM AT JLAB KL2016

FEBRUARY 1-3, 2016
JEFFERSON LAB
NEWPORT NEWS, VIRGINIA

SCOPE

The Workshop is following Lot12-15-001 "Physics Opportunities with Secondary KL beam at JLab" and will be dedicated to the physics of hyperons produced by the kaon beam on unpolarized and polarized targets with GlueX set up in Hall D. The emphasis will be on the hyperon spectroscopy. Such studies could contribute to the existing scientific program on hadron spectroscopy at Jefferson Lab.

The Workshop will also aim at boosting the international collaboration, in particular between the US and EU research institutions and universities.


The Workshop would help to address the comments made by the PAC43, and to prepare the full proposal for the next PAC44.

ORGANIZING COMMITTEE

Moskov Amaryan, ODU, chair
Eugene Chudakov, JLab
Curtis Meyer, CMU
Michael Pennington, JLab
James Ritman, Ruhr-Uni-Bochum & IKP Jülich
Igor Strakovsky, GWU

WWW.JLAB.ORG/CONFERENCES/KL2016





72 people & 28 talks

Doenigus Huovinen Tsuchikawa

YSTAR

Excited Hyperons in QCD Thermodynamics at Freeze-Out

2016

NOVEMBER 16 - 17, 2016

Jefferson Lab
Newport News, Virginia

ORGANIZING COMMITTEE

Moskov Amaryan - Chair James Ritman, Ruhr ODU
U. Babson & IKP Jülich
Eugene Chudakov Igor Strakovsky JLab GWU
Kishana Rajagopal MIT
Chandla Ratti University of Houston

A workshop to discuss the influence of possible "missing" hyperon resonances (JLab KLF Project) on QCD thermodynamics, on freeze-out in heavy ion collisions and in the early universe, and in spectroscopy. Recent studies that compare lattice QCD calculations of thermodynamic calculations, statistical hadron resonance gas models, and ratios between measured yields of different hadron species in heavy ion collisions provide indirect evidence for the presence of "missing" resonances in all of these contexts. The aim of the workshop is to sharpen these comparisons, advance our understanding of the formation of baryons from quarks and gluons microseconds after the Big Bang and in today's experiments, and to connect these developments to experimental searches for direct, spectroscopic, evidence for these resonances. This Workshop is a successor to the recent KL2016 Workshop

WWW.JLAB.ORG/CONFERENCES/YSTAR2016/

THE GEORGE WASHINGTON UNIVERSITY JÜLICH OLD DOMINION UNIVERSITY Jefferson Lab ES

Arriola

Edwards

Xie

Goity

Montgomery

Manley

Crede

Alba

Guo

Stroth

Noumi Tang

Bellwied Ratti

<https://www.jlab.org/conferences/YSTAR2016/>

Speakers: Mai

Chudakov

Garcilazo

Amaryan

Begun

Noronha-Hostler

Myhrer

Ohnishi

Ritman

Capstick



Thank You



*Moskov Amaryan
Yakov Azimov
William Briscoe
Eugene Chudakov
Pavel Degtyarenko
Michael Döring
Alexander Laptev
Ilya Larin
Maxim Mai
Mark Manley
James Ritman
Simon Taylor*

igor@gwu.edu

