



# Simulation and Reconstruction of $K_L + p \rightarrow K_S + p$ and $K_L + p \rightarrow K^+ + n$ using GlueX tracker and KLF software.

---

Vitaly Baturin, ODU, 08/29/2024

## Outlook

1. Simulated Detector performance: examples  $dE/dx$  vs momentum etc.
2. Reconstruction of  $K_L + p$  and  $K^+ + n$  final states  
 $K_L + p \rightarrow K_S + p$  at **low/high** beam momenta  
 $K_L + p \rightarrow K^+ + n$  at **low/high** beam momenta
3. Background from other halls.
4. Conclusion



Identification of final particles.  $K_L + p \rightarrow K_s + p$  and  $K_L + p \rightarrow K^+ + n$

Hadronic decays, lifetimes, and detector dimensions.



$K_s \rightarrow \pi^+ \pi^-$  69.2 %  $c\tau = 2.9$  cm  $\Rightarrow$  **decays mostly inside LH2** target and close to it.  
 $\rightarrow \pi^0 \pi^0$  30.7 %

$K^+ \rightarrow \pi^+ \pi^+ \pi^-$  5.6 %  $c\tau = 371.2$  cm  $\Rightarrow$   **$K^+$  almost "stable"** within the LH2 Target and CDC.  
 $\rightarrow \pi^+ \pi^0$  20.7 %

- $K_s + p$  reconstruction: via  $\pi^+ \pi^-$  tracks; **EM**( $\pi^+ \pi^-$ ) for  $K_s$  and **MM**( $\pi^+ \pi^-$ ) for proton.
- $K^+ + n$  reconstruction: via  $K^+$ -track and **MM**( $K^+$ ) for neutron.
- Using dE/dx in CDC.






Example.  $dE/dx$  in CDC of Gluex Detector .vs. particle momentum for



- Good separation of pions below  $\sim 1.4$  GeV/c and  $\sim 0.7$  GeV/c for  $K^+$ .







# Example. Generated .vs. Reconstructed momenta in $K_L + p \rightarrow K_S(\pi^+\pi^-) + p$ .



- Good reconstruction at all generated momenta.





# Example. Reconstruction of $\pi^+\pi^-$ vertex for $K_L+p \rightarrow K_S(\pi^+\pi^-)+p$ .



- LH2 target sized in cm as  $r \times z = 3 \text{ cm} \times 40 \text{ cm}$ .
- LH2 target is well reproduced by the  $\pi^+\pi^-$  vertex coordinates.
- Diffuse area around  $r=3 \text{ cm}$  is of 1 cm ( $\Rightarrow \sigma \sim 0.25 \text{ cm}$ ).
- Long exponential r-tail is due to  $K_S$  lifetime ( $c\tau=2.9 \text{ cm}$ )







# Example. Reconstruction of $\pi^+\pi^-$ vertex in $K_L+p \rightarrow K_S(\pi^+\pi^-)+p$ . MM( $\pi^+\pi^-$ ) and EM( $\pi^+\pi^-$ ) inside LH2 target.



- EM( $\pi^+\pi^-$ ) resolution is obviously better inside the LH2 target while the background is significantly lower.



# Example. Angular distribution of $\pi^+$ , $\pi^-$ , and $p$ . $K_L + p \rightarrow K_S(\pi^+\pi^-) + p$ .






Reconstruction of  $K_L + p \rightarrow K_S(\pi^+ \pi^-) + p$  at  $K_L$  momentum (0.3,0.6) GeV/c.

Effect of  $\pi^+ \pi^-$  vertex.

## Hadronic decays and lifetimes

$$\begin{aligned} K_S &\rightarrow \pi^+ \pi^- & 69.2 \% & \quad c\tau = 2.9 \text{ cm} \\ &\rightarrow \pi^0 \pi^0 & 30.7 \% & \end{aligned}$$









# Effect of vertex on $K_s(\pi^+\pi^-)$ reconstruction in $K_L + p \rightarrow K_s(\pi^+\pi^-) + p$ at $K_L$ momentum (0.3,0.6) GeV/c.

Inside LH2 target  $r < 3$  cm, DOCA  $< 0.45$  cm




- With **vertex cuts** (left) Rec. Eff. **~25%** (=2500/11000), while the sensitivity (=peak/pedestal)= **~100**.
- No cuts (right) **~50%** (=6000/11000); the sensitivity **~10 times lower**.




# Effect of $\pi^+\pi^-$ vertex on **proton** reconstruction in $K_L+p \rightarrow K_S(\pi^+\pi^-)+p$ at beam momentum (0.3,0.6) GeV/c.

$K_S$ -mass cut, vertex  $r < 3$  cm, DOCA  $< 0.45$  cm.




$K_S$ -mass cut via EM( $\pi^+\pi^-$ ). No vertex cuts.



- Inside LH2 reconstruction efficiency of proton drops from **~44%** to **~22%**.
- Little change of the background, but better MM resolution.





Effect of  $\pi^+ \pi^- (K_S)$  vertex reconstruction  $K_L + p \rightarrow K_S (\pi^+ \pi^-) + p$   
at high  $K_L$  beam momenta (0.55, 4.55) GeV/c.





# Reconstruction of $K_s$ and $p$ in $K_L + p \rightarrow K_s + p$ at $K_L$ momentum (0.55,4.55) GeV/c. Vertex cuts.




- In wide domain of beam momentum Reconstruction Efficiencies are  $\sim 14\%$  for  $K_s$  and  $\sim 2\%$  for protons.




Reconstruction of  $K_s$  and  $p$  in  $K_L + p \rightarrow K_s + p$  at  $K_L$  momentum (0.35,4.55) GeV/c. Effective and Missing mass of  $\pi^+ \pi^-$  pairs vs beam momentum.

LH2 cuts applied






$K_s$  mass and LH2 cuts applied





# Reconstruction of **proton** in $K_L + p \rightarrow K_S (\pi^+ \pi^-) + p$ at **generated $K_L$ energy (0.1,0.3) GeV**



- #tracks >=1 &  $|EM(\pi^+ \pi^-) - m_K| < 20$  MeV. **No vertex cuts**.
- Overall Rec. Efficiency =  $0.83 * 0.63 = \sim 53\%$  for protons and  $\sim 75\%$  for K<sub>s</sub>






# Reconstruction of $K_s+n$ state in

$K_L+p \rightarrow K^+ + n$  at  $K_L$  momentum (0.35,0.55) GeV/c.

Hadronic decay

$K^+ \rightarrow \pi^+ \pi^+ \pi^-$  5.6 %  $c\tau = 371.2$  cm (“stable”)


$\rightarrow \pi^+ \pi^0$  20.7 %






# Reconstruction of neutron in $K_L + p \rightarrow K^+ + n$ with vertex cuts.

$K_L$  momentum (0.35,0.55) GeV/c



$K_L$  momentum (0.55,4.55) GeV/c



- Reconstruction efficiency of neutrons ~50%.







## Backgrounds to $K_L + p \rightarrow K^+ + n$

- (1)  $K_L + p \rightarrow \pi^+ + \Sigma^0(\Lambda)$ ;  $p_{\text{Kaon}} > 0$
- (2)  $n + p \rightarrow K_s + \Sigma^+ + n$ ;  $p_{\text{neutron}} > 2.6 \text{ GeV}/c \Rightarrow \text{low } n\text{-flux.}$
- (3) Beam leak from other halls and reconstruction of  $K_L + p \rightarrow K^+ + n$




# Background reaction $K_L + p \rightarrow \pi^+ + \Sigma^0$



- Well separated from proton and neutron MM peaks at low beam momenta.





# $K_L + p \rightarrow K^+ + \Xi^0$ at $0.35 < p_K < 0.55$ GeV/c . Missing Mass of $K^+$ and beam leak from other halls (left) vs normal Hall D beam (right)



- No structures in Missing Mass of  $K^+$  !
- At  $p_{\text{beam}}=0.45$  GeV/c the “leak” / “beam” ratio =  $\sim 100/10 = \sim 10$ ; we expect  $\sim 2$  (factor 5).
- The ratio of neutron\_peak to leak of “ $\Xi^0$ ” =  $\sim 300/75 = \sim 4$ ; we expect  $\sim 5 \cdot 4 = 20$ , i.e.  **$\sim 5\%$  background.**



# $K_L + p \rightarrow K^+ + \Xi^0$ . Background oscillation caused by beam leak.




- The background part shows oscillations (10% amplitude) due to beam leak.



# What to do with beam leak background?

1. FLUKA shows that **minum energy of  $K_L$**  at LH2 target  $T=0.1$  GeV.  
 $\Rightarrow K_L$  **momentum**  $p_K = 0.33$  GeV/c  
 $\Rightarrow \beta_K = p_K / (T + m_K) = 0.33 / 0.597 = 0.554$   
 $TOF_K = 2400[\text{cm}] / (30[\text{cm/ns}] * 0.554) = \mathbf{144}$  ns  
 $TOF_\gamma = 2400[\text{cm}] / (30[\text{cm/ns}] * 1.000) = \mathbf{80}$  ns
2. The difference  $TOF_K - TOF_\gamma = \mathbf{64}$  ns , therefore  
all **beam  $K_L$** 's with  $p > 0.33$  GeV/c do **fit into (0,64) ns** interval!
3. **Provided 128 ns** between bunches the following (64,128) ns interval  
is **filled by beam leak** only. Therefore this interval may be used  
to **permanently measure /subtract the background** .





## Conclusion

Final state (FS)	#FS/#K <sub>L</sub> / Resol.	#FS/#K <sub>L</sub> / Resol.
K <sub>L</sub> beam mom.	0.3-0.6 GeV/c	0.5-5.0 GeV/c
→K <sub>s</sub> + ...	53 % / 10 MeV	14 % / 20 MeV
→K <sub>s</sub> +p	44 % / 20 MeV	2 % / 25 MeV
→K <sup>+</sup> +n	50 % / 15 MeV	6 % / 50 MeV
→π <sup>+</sup> +Σ <sup>0</sup>	64 % / 10 MeV	18 % / 15 MeV

- π<sup>+</sup>π<sup>-</sup>
- GlueX CDC is an **ideal detector** at K<sub>L</sub> beam momentum (0.3, 0.6) GeV/c.
- Overage **reconstruction efficiency** ~50% in this region.
- **Advantage** of K<sub>L</sub>+p → K<sub>s</sub>(π<sup>+</sup>π<sup>-</sup>)+p is that it has **3 charged particles** of low momenta, hence - better **resolution** and **vertex** localisation; good cross check for K<sub>L</sub>+p → K<sup>+</sup>+n.
- **Beam leak** background **does not create problems** for neutron rec. via MM(K<sup>+</sup>).