

Proposal for JLab PAC46

***Strange Hadron Spectroscopy with
Secondary K_L Beam at GlueX***

Moskov Amaryan

(For KLF Collaboration)

JLab Management Briefing, May 8, 2018

Outline

Physics Motivation

- *Hyperon Spectroscopy*
- *Strange Meson Spectroscopy*

K_L Facility at JLab

- *Electron Beam*
- *Compact Photon Source*
- *Be Target*
- *Flux Monitor*
- *K_L Beam*
- *LH_2/LD_2 Target*

Summary



Hyperon Spectroscopy

According *LQCD* there should be

many more states including hybrids (thick bordered)

8-states

5-states





6-states

4-states

3-states

4-states





1-state



1-states

Proposed Measurements on Proton Target

existing data





KLF 100 days



Proposed Measurements on Neutron Target

Sensitivity to different solutions



100 days on LD_2 target

Search for Hyperon Resonances with PWA


For Scattering experiments on both proton & neutron targets we need to determine:

- differential cross sections &**
- self polarization of strange hyperons**
- perform coupled-channel PWA**
- look for poles in complex energy plane
(not naïve bump hunting)**
- identify Λ^* , Σ^* , Ξ^* & Ω^* up to 2400 MeV**


As kaon nucleon scattering data are very poor

We use pion nucleon scattering data with statistics generated according to expected KLF data for 20(100) days to show PWA sensitivity to obtain results close to the best fit

Using πp Scattering




Statistics was generated according to KLF for



Obviously: we need **at least 100 days** to get unique solution

Strange Meson Spectroscopy


$K\pi$ Scattering



Proposed Measurements


SLAC

$$K^- \pi^+ \rightarrow K^- \pi^+$$




Belle

$$\tau \rightarrow K \pi \nu_\tau$$



KLF

$$K_L \pi^0 \rightarrow K^+ \pi^-$$





region of $\mathcal{K}(800)$


Proposed Measurement

I=1/2

S-Wave



Scalar Meson Nonet




Four states called κ

still need further confirmation(PDG)


We can measure all of them

Measurement of $\kappa(800)$



100 days of running

Hall-D beamline and GlueX Setup



Electron Beam Parameters

$$E_e = 12 \text{ GeV} \quad I = 5 \mu\text{A}$$

$$\text{Repetition rate} \quad 64 \text{ ns}$$

No major problems.

Doable !


Confirmed by Todd Satogata

Estimated investment \$60 K

Compact Photon Source

Conceptual design is completed for Halls C/A
for $I = 2.7 \mu A$


Could be extended for $I = 5 \mu A$
in Hall D



Meets RadCon Radiation Requirements

Estimated cost \$1.5-2.0 M

Be Target Assembly: Conceptual Design





-Meets RadCon Radiation Requirements

-Conceptual Design Endorsed by Hall-D Engineering Staff

Estimated cost ~\$1.2 M

Flux Monitor



Reconstructed K_L mass


Flux measurement star. err. <1%

Estimated syst. err. <3%

Pursuing UK funding


Estimated cost ~\$ 0.7 M

K_L Beam Flux




$$N(K_L)/sec \sim 10^4$$

SLAC




$$\frac{N(K_L)_{JLAB}}{N(K_L)_{SLAC}} \sim 10^3$$



LH₂/LD₂ Cryogenic Target for Neutral Kaon Beam at Hall D

The GlueX liquid hydrogen target.



Current

Proposed & Feasible


Longer and ticket target is needed to enhance production rate

Conceptual design endorsed by target group (Chris Keith)

Estimated cost ~\$ 30 K


$K\pi$ Scattering Resolutions

Four Momentum Resolution for $K_L p \rightarrow K^+ \pi^- p$



-Good resolution at low- t is needed to be on pion pole

$K^+ \pi^-$ Invariant Mass Resolution for $K_L p \rightarrow K^+ \pi^- p$



-Binning in $\sim 10 \text{ MeV}$ will cover almost entire elastic K - π scattering range

PHYSICS WITH NEUTRAL KAON BEAM AT JLAB KL2016

FEBRUARY 1-3, 2016
JEFFERSON LAB
NEWPORT NEWS, VIRGINIA

SCOPE

The Workshop is following L012-15-001 "Physics Opportunities with Secondary KL beam at JLab" and will be dedicated to the physics of hyperons produced by the kaon beam on unpolarized and polarized targets with GlueX set up in Hall D. The emphasis will be on the hyperon spectroscopy. Such studies could contribute to the existing scientific program on hadron spectroscopy at Jefferson Lab.

The Workshop will also aim at boosting the international collaboration, in particular between the US and EU research institutions and universities.

The Workshop would help to address the comments made by the PAC43, and to prepare the full proposal for the next PAC44.

ORGANIZING COMMITTEE

Moskov Amaryan, ODU, chair
Eugene Chudakov, JLab
Curtis Meyer, CMU
Michael Pennington, JLab
James Ritman, Ruhr-Uni-Bochum & IKP Jülich
Igor Strakovsky, GWU

WWW.JLAB.ORG/CONFERENCES/KL2016



YSTAR Excited Hyperons in QCD Thermodynamics at Freeze-Out 2016

NOVEMBER 16-17, 2016

Jefferson Lab
Newport News, Virginia

A workshop to discuss the influence of possible "missing" hyperon resonances (JLab KLF Project) on QCD thermodynamics, on freeze-out in heavy ion collisions and in the early universe, and in spectroscopy. Recent studies that compare lattice QCD calculations of thermodynamic calculations, statistical hadron resonance gas models, and ratios between measured yields of different hadron species in heavy ion collisions provide indirect evidence for the presence of "missing" resonances in all of these contexts. The aim of the workshop is to sharpen these comparisons, advance our understanding of the formation of baryons from quarks and gluons microseconds after the Big Bang and in today's experiments, and to connect these developments to experimental searches for direct, spectroscopic, evidence for these resonances. This Workshop is a successor to the recent KL2016 Workshop

ORGANIZING COMMITTEE

Moskov Amaryan - Chair, ODU
Eugene Chudakov, JLab
Krishna Rajagopal, MIT
Claudia Ratti, University of Houston
James Ritman, Ruhr U. Bochum & IKP Jülich
Igor Strakovsky, GWU



WWW.JLAB.ORG/CONFERENCES/YSTAR2016/




HIPS 2017

New Opportunities with High-Intensity Photon Sources

February 6-7, 2017
Catholic University of America
Washington, DC U.S.A.

This workshop aims at producing an optimized photon source concept with potential increase of scientific output at Jefferson Lab, and at refining the science for hadron physics experiments benefiting from such a high-intensity photon source. The workshop is dedicated to bringing together the communities directly using such sources for photo-production experiments, or for conversion into K_s beams. The combination of high precision calorimetry and high intensity photon sources can provide greatly enhanced scientific benefit to (deep) exclusive processes like wide-angle and time-like Compton scattering. Potential prospects of such a high-intensity source with modern polarized targets will also be discussed. The availability of K_s beams would open new avenues for hadron spectroscopy, for example for the investigations of "missing" hyperon resonances, with potential impact on QCD thermodynamics and on freeze-out both in heavy ion collisions and the early universe.



Organizing Committee:

Tanja Horn - CUA
Cynthia Keppel - JLab
Carlos Munoz-Camacho - IPNO
Igor Strakovsky - GWU



π-K Interactions Workshop





ORGANIZING COMMITTEE

Moskov Amaryan, ODU (Chair)
U.-G. Meissner, U. Bonn/FZ Jülich
Curtis Meyer, CMU
James Ritman, Ruhr-Uni-Bochum/FZ Jülich
Igor Strakovsky, GWU

February 14-15, 2018

Jefferson Lab • Newport News, VA

The π-K scattering enables direct investigations of scalar and vector K* states, including the not yet established S-wave k(800) state. These studies are also needed to get precise values of vector and scalar form factors: to independently extract CKM matrix element V_{us} and to test the Standard Model unitarity relation in the first row of CKM matrix, to study CP violation from the Dalitz plot analysis of open charm D meson decays and in a charmless decays of B mesons in K_s final states. Significant progress is made lately in Lattice QCD, in the phenomenology and in the Chiral Perturbation Theory to describe different aspects of π-K scattering. The main source of experimental data is based on experiments performed in SLAC almost five decades ago at 1970-80s. The recently proposed KL Facility incorporating the GlueX spectrometer at JLab will be able to improve the π-K scattering database by about three orders of magnitude in statistics. The workshop will discuss the necessity for and the impact of the future high statistics data obtained at JLab on π-K scattering.



<https://www.jlab.org/conferences/pki2018/>

KL2016

[60 people from 10 countries, 30 talks] <https://www.jlab.org/conferences/kl2016/>

OC: M. Amaryan, E. Chudakov, C. Meyer, M. Pennington, J. Ritman, & I. Strakovsky

YSTAR2016

[71 people from 11 countries, 27 talks] <https://www.jlab.org/conferences/YSTAR2016/>

OC: M. Amaryan, E. Chudakov, K. Rajagopal, C. Ratti, J. Ritman, & I. Strakovsky

HIPS2017

[43 people from 4 countries, 19 talks] <https://www.jlab.org/conferences/HIPS2017/>

OC: T. Horn, C. Keppel, C. Munoz-Camacho, & I. Strakovsky

PKI2018

[48 people from 9 countries, 27 talks] <http://www.jlab.org/conferences/pki2018/>

OC: M. Amaryan, U.-G. Meissner, C. Meyer, J. Ritman, & I. Strakovsky

In total: 222 participants & 103 talks

Proposal:
200 Members
61 Institutions
20 Countries

A. Ali¹⁸, M. B. Ali⁴⁷, M. J. Amaryan^{44,*†}, E. G. Anassontzis², A. V. Anisovich^{4,48},
A. Austregesilo³⁰, M. Baalouch⁴⁵, F. Barbosa³⁰, J. Barlow¹³, A. Barnes⁷, E. Barriga¹³,
M. Bashkanov^{10,†}, A. Bazavov³⁹, T. D. Beattie⁵⁰, R. Bellwied²⁰, V. V. Berdnikov⁸, T. Black⁴²,
W. Boeglin¹², M. Boer⁸, W. J. Briscoe¹⁴, T. Britton³⁰, W. K. Brooks⁵³, B. E. Cannon¹³, N. Cao²²,
E. Chudakov³⁰, P. L. Cole²¹, S. Cole¹, V. Crede¹³, M. M. Dalton³⁰, T. Daniels⁴², D. Day⁵⁸,
P. Degtyarenko³⁰, A. Deur³⁰, S. Dobbs¹³, G. Dodge⁴⁵, A. G. Dolgolenko²⁷, M. Döring^{14,30},
M. Dugger¹, R. Dzhygadlo¹⁸, S. Eidelman^{5,44}, R. Edwards³⁰, H. Egiyan³⁰, A. Ernst¹³,
A. Eskandarian¹⁴, P. Eugenio¹³, C. Fanelli³⁶, S. Fegan¹⁴, A. Filippi²⁵, A. M. Foda⁵⁰, J. Frye²³,
S. Furletov³⁰, L. Gan⁴², A. Gasparyan⁴¹, G. Gavalian³⁰, M. Gauzshtein^{54,55}, N. Gevorgyan⁶¹,
C. Gleason²³, D. I. Glazier¹⁷, J. Goity^{30,19}, V. S. Goryachev²⁷, K. Götzen¹⁸, A. Goncalves¹³,
L. Guo¹², H. Haberzettl¹⁴, M. Hadžimehmedović⁵⁷, H. Hakobyan⁵³, A. Hamdi¹⁸, S. Han⁶⁰,
J. Hardin³⁶, A. Hayrapetyan¹⁶, G. M. Huber⁵⁰, A. Hurley⁵⁹, C. E. Hyde⁴⁵, T. Horn⁸,
D. G. Ireland¹⁷, M. Ito³⁰, N. Jarvis⁷, R. T. Jones⁹, V. Kakoyan⁶¹, G. Kalicy⁸, M. Kamel¹²,
C. D. Keith³⁰, C. W. Kim¹⁴, F. J. Klein¹⁴, C. Kourkoumeli², G. Krafft³⁰, S. Kuleshov⁵³,
I. Kuznetsov^{54,55}, A. B. Laptev³³, I. Larin³⁵, D. Lawrence³⁰, D. I. Lersch¹³, M. Levillain⁴¹, H. Li⁷,
W. Li⁵⁹, K. Livingston¹⁷, B. Liu²², G. J. Lolos⁵⁰, V. E. Lyubovitskij^{56,54,55,53}, D. Mack³⁰,
M. Mai¹⁴, D. M. Manley³¹, M. Mazouz⁴⁷, H. Marukyan⁶¹, V. Mathieu³⁰, P. T. Mattione³⁰,
M. Matveev⁴⁸, V. Matveev²⁷, M. McCaughan³⁰, W. McGinley⁷, M. McCracken⁷, J. McIntyre⁹,
U.-G. Meißner^{4,29}, V. Mokeev³⁰, F. Nerling¹⁸, C. A. Meyer⁷, R. Miskimen³⁵, R. E. Mitchell²³,
F. Mokaya⁹, C. Morningstar⁷, B. Moussallam⁴⁶, K. Nakayama¹⁵, Y. Oh³², R. Omerović⁵⁷,
H. Osmanović⁵⁷, A. Ostrovidov¹³, Z. Papandreou⁵⁰, K. Park³⁰, E. Pasyuk³⁰, M. Patsyuk³⁶,
P. Pauli¹⁷, R. Pedroni⁴¹, J. R. Pelaez³⁴, L. Pentchev³⁰, M. R. Pennington¹⁷, K. J. Peters¹⁸,
W. Phelps¹⁴, A. Pilloni³⁰, E. Pooser³⁰, J. W. Price⁶, N. Qin⁴³, J. Reinhold¹², D. Richards³⁰,
D.-O. Riska¹¹, B. Ritchie¹, J. Ritman^{51,28,†}, L. Robison⁴³, A. Rodas³⁴, D. Romanov³⁷,
C. Romero⁵³, J. Ruiz de Elvira³, H.-Y. Ryu⁴⁹, C. Salgado⁴⁰, E. Santopinto²⁴, A. V. Sarantsev^{4,48},
T. Satogata³⁰, A. Schertz⁵⁹, R. A. Schumacher⁷, C. Schwarz¹⁸, J. Schwiening¹⁸,
A. Yu. Semenov⁵⁰, I. A. Semenova⁵⁰, K. K. Seth⁴³, X. Shen²², M. R. Shepherd²³, E. S. Smith³⁰,
D. I. Sober⁸, D. Sokhan¹⁷, A. Somov³⁰, S. Somov³⁷, O. Soto⁵³, M. Staib⁷, J. Stahov⁵⁷,
J. R. Stevens^{59,†}, I. I. Strakovsky^{14,†}, A. Švarc⁵², A. Szczepaniak^{23,30}, V. Tarasov²⁷, S. Taylor³⁰,
A. Teymurazyan⁵⁰, A. Trabelsi⁴⁷, G. Vasileiadis², D. Watts¹⁰, D. Werthmüller¹⁷, T. Whitlatch³⁰,
N. Wickramaarachchi⁴⁵, M. Williams³⁶, B. Wojtsekhowski³⁰, R. L. Workman¹⁴, T. Xiao⁴³,
Y. Yang³⁶, N. Zachariou¹⁰, J. Zarling²³, J. Zhang⁵⁸, Z. Zhang⁶⁰, G. Zhao²², B. Zou²⁶, Q. Zhou²²,
X. Zhou⁶⁰, B. Zihlmann³⁰

SUMMARY

-Proposed KL Facility has unique capability to improve existing world database up to three orders of magnitude

-In Hyperon spectroscopy

PWA will allow to measure pole positions and widths of excited hyperon states

-In Strange Meson Spectroscopy

PWA will allow to measure excited K^* states including scalar $f_0(800)$ states

- To accomplish physics program 100 days per LH2 and LD2 is required

-All components of KL Facility considered are feasible

-With total cost less than \$ 4.0 M

CPS(~\$ 1.5-2.0 M)

BeTarget(~\$ 1.2 M)

FluxMonitor (~\$ 0.7 M)

Electron Beam (~\$ 60K)



Cryo-Targets (~\$ 30K)

Backup

Possible Other Impacts

$$Br(K_L \rightarrow \pi e \nu) \sim |f_+(0) V_{us}|^2$$

$$Br(\tau \rightarrow K \pi \nu) \sim |f_+(0) V_{us}|^2$$



FLAG Collaboration Eur.Phys.J. C77 (2017) no.2, 112

Eur.Phys.J. C69 (2010) 399-424

No precise data yet !

test of unitarity:

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$$