

Highlights for KLF — a personal view

Eberhard Klempf

University of Bonn



KLF Collaboration Meeting

Virtual meeting online

December 9, 2020

Introduction

► What does QCD teach us:

Lattice QCD	Quark model	Dynamically gen. states
QM states + glueballs and hybrids	not all dynamically generated states are QM states	not all QM states are dynamically generated

Is this a change of the quantum mechanical basis?
 $(qqq \quad \text{versus} \quad \text{baryon +meson})$

The relation between QM states and dynamically generated states is not understood

QCD inspired models predict states beyond the quark model!

1. Baryons in the quark model

SU(6) relates the spectrum of Λ and Σ hyperons to N and Δ :

$$6 \otimes 6 \otimes 6 = 56_S \quad \oplus \quad 70_M \oplus 70_M \quad \oplus \quad 20_A.$$

$$56 = {}^4\text{10} \oplus {}^2\text{8}, \quad 70 = {}^2\text{10} \oplus {}^4\text{8} \oplus {}^2\text{8} \oplus {}^2\text{1} \quad 20 = {}^2\text{8} \oplus {}^4\text{1}.$$

Δ and N Δ and N and Λ N

Σ & Σ, Λ Σ and Σ, Λ Σ, Λ and Λ

The number of Σ resonances is equal to $n_\Sigma = n_N + n_\Delta$.


The number of Λ resonances is equal to $n_\Lambda = n_N + n_{\Lambda_8}$.

→ The Σ and Λ spectra are very rich!


- ▶ What do we know?
- ▶ Can we identify singlet, octet and decuplet states?
- ▶ There should be spin-quartet Λ singlet states!

1.1 Large number of missing resonances!

More data needed! Prime goal of KLF.



Prominent features: $\Sigma(1775)5/2^-$



$\Lambda(1520)3/2^-$ and $\Lambda(1820)5/2^+$

1.2 Baryons have two oscillators: Understand first and second excitation shell!

The negative-parity states:

$(D, L_N^P) S J^P$	Singlet	Octet			Decuplet	
$(70, 1^-_1) \frac{1}{2} \frac{1}{2}^-$	$\Lambda(1380)$	$N(1535)$	$\Lambda(1670)$ +135	$\Sigma(1620)$ +85	$\Delta(1620)$	$\Sigma(1900)\dagger$
	$\Lambda(1405)$	$N(1520)$	$\Lambda(1690)$ +170	$\Sigma(1670)$ +150	$\Delta(1700)$	$\Sigma(1910)\dagger$
	$\Lambda(1520)$	$N(1650)$	$\Lambda(1800)$ +150	$\Sigma(1750)$ +100		
		$N(1700)$	-	-		
		$N(1675)$	$\Lambda(1830)$ +155	$\Sigma(1775)$ +100	\dagger 1800 and 2050 MeV expected	
		Mean:	+150 MeV	+110 MeV		

1st excitation shell: single-oscillator excitations!

2nd excitation shell: 56plet — single-oscillator excitations!

70plet — mixed !

20plet — two-oscillator excitations!

Number of expected and observed resonances that can be assigned to the 2nd excitation shell for $J^P = 1/2^+, .., 7/2^+$. The first number gives the expected number of resonances, followed by the number of observed resonances with 3* and 4*, 1* and 2* (in parentheses).

		1/2 ⁺	3/2 ⁺	5/2 ⁺	7/2 ⁺	Sum
N	expected (4*3*, 2*1*):	4 (4,0)	5 (3,1)	3 (1,2)	1 (1,0)	13 (9,3)
Δ		2 (1,1)	3 (2,0)	2 (1,0)	1 (1,0)	8 (5,1)
Λ		6 (2,1)	7 (1,1)	5 (2,0)	1 (0,1)	19 (5,3)
Σ		6 (1,1)	8 (0,4)	5 (1,1)	2 (1,0)	21 (3,6)
Ξ		6 (0,0)	8 (0,0)	5 (0,0)	2 (0,0)	21 (0,0)
Ω		2 (0,0)	3 (0,0)	2 (0,0)	1 (0,0)	8 (0,0)

In the spectra of Λ and Σ resonances, 40 resonances are expected in the second excitation shell. Only 8 are known with 3 or 4 stars in RPP notation.

That is too little!

$(D, L_N^P) S J^P$	Singlet	Octet			Decuplet		
$(56, 0_2^+)$ $\frac{1}{2} \frac{1}{2}^+$ $\frac{3}{2} \frac{3}{2}^+$ $\frac{5}{2} \frac{5}{2}$		$N(1440)$		$\Lambda(1600)$	$\Sigma(1660)$	$\Delta(1600)$	$\Sigma(1780)$
$(70, 0_2^+)$ $\frac{1}{2} \frac{1}{2}^+$ $\frac{3}{2} \frac{3}{2}^+$ $\frac{5}{2} \frac{5}{2}$	$\Lambda(1710)$	$N(1710)$		$\Lambda(1810)$	$\Sigma(1880)$	$\Delta(1750)$	-
$(56, 2_2^+)$ $\frac{1}{2} \frac{3}{2}^+$ $\frac{1}{2} \frac{5}{2}^+$ $\frac{3}{2} \frac{1}{2}^+$ $\frac{3}{2} \frac{3}{2}^+$ $\frac{3}{2} \frac{5}{2}^+$ $\frac{3}{2} \frac{7}{2}^+$		$N(1720)$		$\Lambda(1890)$	$\Sigma(1940)$	$\Delta(1910)$	-
		$N(1680)$		$\Lambda(1820)$	$\Sigma(1915)$	$\Delta(1920)$	$\Sigma(2080)$
						$\Delta(1905)$	$\Sigma(2070)$
						$\Delta(1950)$	$\Sigma(2030)$
$(70, 2_2^+)$ $\frac{1}{2} \frac{3}{2}^+$ $\frac{5}{2}^+$ $\frac{3}{2} \frac{1}{2}^+$ $\frac{3}{2} \frac{3}{2}^+$ $\frac{5}{2}^+$ $\frac{7}{2}^+$	$\Lambda(2070)$ $\Lambda(2110)$ $(\text{all } 1^* \text{ in RPP})$	$-$		$-$	$-$	$-$	-
		$N(1860)$		$-$	$-$	$\Delta(2000)$	-
		$N(1880)$		$-$	$-$		
		$N(1900)$		$-$	$-$		
		$N(2000)$		$-$	$-$		
		$N(1990)$		$\Lambda(2085)$	$-$		
$(20, 1_2^+)$ $\frac{1}{2} \frac{1}{2}^+$ $\frac{3}{2}^+$ $\frac{5}{2}^+$	$-$ $-$ $-$	$-$		$-$	$-$	Not yet any two-oscillator excitation	

1.3 Search for the 20plet:

$$\Lambda(2099)1/2^+, \Lambda(2176)3/2^+, \Lambda(2150)5/2^+$$

U. Löring, B. C. Metsch and H. R. Petry, Eur. Phys. J. A 10 447 (2001).

The multiplets 70, 56, and 20 arise from the combination of the three light quarks u, d, s having spin 1/2:

$$6 \otimes 6 \otimes 6 = 56_S \oplus 70_M + 20_A$$


2nd excitation shell:

$$S = \frac{1}{\sqrt{2}} \{ [\phi_{0s}(\vec{\rho}) \times \phi_{0d}(\vec{\lambda})] + [\phi_{0d}(\vec{\rho}) \times \phi_{0s}(\vec{\lambda})] \}^{(L=2)},$$


$$M_S = \frac{1}{\sqrt{2}} \{ [\phi_{0s}(\vec{\rho}) \times \phi_{0d}(\vec{\lambda})] - [\phi_{0d}(\vec{\rho}) \times \phi_{0s}(\vec{\lambda})] \}^{(L=2)}$$

$$M_A = [\phi_{0p}(\vec{\rho}) \times \phi_{0p}(\vec{\lambda})]^{(L=2)},$$

$$A = [\phi_{0p}(\vec{\rho}) \times \phi_{0p}(\vec{\lambda})]^{(L=1)}.$$



Classical orbits:



Baryons in the 20plet can neither decay nor be produced in a single step!

A. Thiel et al. [CBELSA/TAPS], Phys. Rev. Lett. 114 091803 (2015).

Search for 20plet- Λ resonances in a cascade process:

$$\Sigma^+(\text{highmass}) \rightarrow \Lambda(2099)1/2^+ + \pi^+ \quad \begin{matrix} & & \text{to singlet component of } \eta \text{ only} \\ & & \uparrow \end{matrix}$$

$$\downarrow \rightarrow \Lambda(1520)3/2^- + \eta, L = 0$$
$$\downarrow \rightarrow \Sigma + \pi, N\bar{K}$$

$$\Sigma^+(\text{highmass}) \rightarrow \Lambda(2176)3/2^+ + \pi^+$$

$$\downarrow \rightarrow \Lambda(1820)5/2^+ + \pi, L = 0$$
$$\downarrow \rightarrow \Sigma + \pi, N\bar{K}$$

$$\Sigma^+(\text{highmass}) \rightarrow \Lambda(2150)5/2^+ + \pi^+ \quad \begin{matrix} & & \text{to singlet component of } \eta \text{ only} \\ & & \uparrow \end{matrix}$$

$$\downarrow \rightarrow \Lambda(1520)3/2^- + \eta, L = 0$$
$$\downarrow \rightarrow \Sigma + \pi, N\bar{K}$$

Small cross section expected! Questionable due to low expected rate,
but clear proof of three-body dynamics in baryons

2. Dynamically generated resonances beyond the q.m.

2.1 DGR from EFTs

Effective field theories generate
in $\bar{K}N - \Sigma\pi$ coupled
channel dynamics two poles
in the $\Lambda(1405)$ region!


I=0: (1379.2 - i 27.6) MeV
(1433.7 - i 11.0) MeV

J. A. Oller and U. G. Meissner, Phys. Lett. B 500, 263 (2001),
D. Jido, J. A. Oller, E. Oset, A. Ramos, U. G. Meissner,
Nucl. Phys. A 725, 181 (2003).

In the quark model (and so far on
the lattice¹), only one singlet
pole is expected at this mass!

¹: R. Pavao, P. Gubler, P. Fernandez-Soler, J. Nieves, M. Oka
and T. T. Takahashi, [arXiv:2010.01270 [hep-lat]].

Is $\Lambda(1405)$ a SU(3) singlet or octet state?



(D, L_N^P) S J^P	QM	EFT
$(70, 1_1^-)$ $\frac{1}{2} \frac{1}{2}^-$	singlet	$\Lambda(1405)$
$(70, 1_1^-)$ $\frac{1}{2} \frac{1}{2}^-$	octet	$\Lambda(1670)$
$(70, 1_1^-)$ $\frac{3}{2} \frac{1}{2}^-$	octet	$\Lambda(1800)$

2.2 SU(3) nature of $\Lambda(1405)$ from its decays

SU(3) coupling constants for hyperon decays and the SU(6) predictions for the coefficient α in decays of octet hyperons.


V. Guzey and M. V. Polyakov, [arXiv:hep-ph/0512355 [hep-ph]].

$$\text{RPP} \quad \begin{aligned} & 8_1 \rightarrow 8 \otimes 8 \\ \begin{pmatrix} N \\ \Sigma \\ \Lambda \\ \Xi \end{pmatrix} \rightarrow \begin{pmatrix} N\pi & N\eta & \Sigma K & \Lambda K \\ N\bar{K} & \Sigma\pi & \Lambda\pi & \Sigma\eta & \Xi K \\ N\bar{K} & \Sigma\pi & \Lambda\eta & \Xi K \\ \Sigma\bar{K} & \Lambda\bar{K} & \Xi\pi & \Xi\eta \end{pmatrix} = \frac{1}{\sqrt{20}} \begin{pmatrix} 9 & -1 & -9 & -1 \\ -6 & 0 & 4 & 4 \\ 2 & -12 & -4 & -2 \\ 9 & -1 & -9 & -1 \end{pmatrix}^{1/2} \\ & 8_2 \rightarrow 8 \otimes 8 \\ \begin{pmatrix} N \\ \Sigma \\ \Lambda \\ \Xi \end{pmatrix} \rightarrow \begin{pmatrix} N\pi & N\eta & \Sigma K & \Lambda K \\ N\bar{K} & \Sigma\pi & \Lambda\pi & \Sigma\eta & \Xi K \\ N\bar{K} & \Sigma\pi & \Lambda\eta & \Xi K \\ \Sigma\bar{K} & \Lambda\bar{K} & \Xi\pi & \Xi\eta \end{pmatrix} = \frac{1}{\sqrt{12}} \begin{pmatrix} 3 & 3 & 3 & -3 \\ 2 & 8 & 0 & 0 \\ 6 & 0 & 0 & 6 \\ 3 & 3 & 3 & -3 \end{pmatrix}^{1/2} \end{aligned}$$


Decay mode	$8 \rightarrow 8 + 8$	$1 \rightarrow 8 + 8$
$\Lambda \rightarrow N\bar{K}$	$\sqrt{\frac{2}{3}}(2\alpha + 1)A_8$	$\frac{1}{2}A_1$
$\Lambda \rightarrow \Sigma\pi$	$2(\alpha - 1)A_8$	$\sqrt{\frac{3}{2}}A_1$
	$^28[56]$	$^28[70]$
α	$\frac{2}{5}$	$\frac{5}{8}$
	$^21[70]$	$^28[56]$
	$\frac{A(\Lambda \rightarrow N\bar{K})}{A(\Lambda \rightarrow \Sigma\pi)}$	$^28[70]$
Sign	\oplus	$-$
	\ominus	$^48[70]$

2.3 BnGa fit to a large data set

- ▶ $K^- N$ scattering
- ▶ $\Delta E + i\frac{\Gamma}{2}$ of $\pi^- p$ atom
- ▶ CLAS data on $\gamma p \rightarrow K^+ (\Sigma^\pm \pi^\mp)$
- ▶ CLAS data on $(\Sigma^0 \pi^0)$ predicted!




Fit with one pole at 1405 MeV:



Fit with one pole: $\Lambda(1405)$ is SU(3) singlet. Fit with two poles: $\bar{K}N$ amplitude flips sign: $\Lambda(1405)$ become SU(3) octet!

3.4 KLF determines SU(3) structure of $\Lambda(1405)$



$\Sigma^+\pi^-$ mass from $K^-p \rightarrow \pi^-\pi^+\pi^\mp\Sigma^\pm$
for events with $M_{\pi^+\pi^\pm\Sigma^\mp}$ compatible with
 $\Sigma(1670)3/2^-$. Number of events /10 MeV.

R. J. Hemingway, Nucl. Phys. B 253, 742 (1985).

BnGa fit: 75% $\Lambda(1405)1/2^-$;
25% $\Sigma(1385)3/2^+$

The signs of the SU(6) amplitudes for


$\Sigma^+(1670)3/2^- \rightarrow \pi^+\Lambda(1405)$;
 $\Lambda(1405) \rightarrow \Sigma^\pm\pi^\mp$ and

$\Sigma^+(1670)3/2^- \rightarrow \pi^+\Sigma(1385)$;
 $\Sigma(1385) \rightarrow \Sigma^\pm\pi^\mp$.

$\Lambda(1405)$ SU(3) structure:		1	8
$\Sigma^+(1670)3/2^-$	\rightarrow	$\Lambda(1405)\pi^+$	+
	\hookrightarrow	$\Sigma^\pm\pi^\mp$	+
Sign of transition amplitude at pole:		\oplus	\ominus
$\Sigma^+(1670)3/2^-$	\rightarrow	$\Sigma^0(1385)\pi^+$	+
	\hookrightarrow	$\Sigma^\pm\pi^\mp$	+
Sign of transition amplitude at pole:		\oplus	\oplus

3. Hybrids and glueballs


3.1 Σ hybrids at KLF




Most promising:

- ▶ Spring breaking leads to orbital angular momentum in the final state
- ▶ $\Lambda(1520)3/2^-$ dominant structure
- ▶ $\Sigma_h^+(2250)3/2^+ \rightarrow \Lambda(1520)3/2^- + \pi^+$ S-wave decay
- ▶ Two $\Sigma_h^+(2250)3/2^+$ and $\Sigma_h^+(2350)3/2^+$ isolated (lattice prediction)

3.2 No glueballs at KLF , but scalar K_0^*




$f_0(980)$	$a_0(980)$	$K_0^*(700)$
$f_0(1500)$	$a_0(1450)$	$K_0^*(1450)$
$f_0(1710)$?
$f_0(2100)$	$a_0(2020)$	$K_0^*(1950)$



Scalar mesons with strangeness, K_0^* series very important.

Existence or not of a $K_0^*(1680)$ helps to decide if $f_0(1710)$ is a glueball!

4. Pentaquarks





The antidecuplet and its quark model decomposition. The antidecuplet predicted by the chiral soliton model describes nucleons in terms of the pion field and not by the number of quarks. The three corner-states are incompatible with a qqq assignment.

Study reactions:

$K_L p \rightarrow \Theta^+(1530) \rightarrow K^+ n$ charge exchange (phase shift!)

$K_L p \rightarrow K_S \Xi^+(2070)$



Summary

My own expectation

KLF offers a wide range of opportunities:

1. Baryons in the quark model

- 1.1 Missing resonances
- 1.2 Two oscillators
- 1.3 Λ resonances in the 20plet

Several new resonances will be found
and/or upgraded

Unlikely, not sufficient statistics

2. Dynamically generated resonances beyond the q.m.

- 2.1 DGR from EFTs
- 2.2 SU(3) nature of $\Lambda(1405)$
- 2.3 The BnGa fit
- 2.4 KLF and the $\Lambda(1405)$

KLF will contribute to clarification

3. Glueballs and hybrids

- 3.1 Σ hybrids at KLF
- 3.2 No glueballs at KLF but scalar K_0^*

“Candidates” will be discussed
Better $K_0^*(700)$, $K_0^*(1450)$, ... ?

4. Pentaquarks

Good new upper limits