

# *K-Long Facility at Jefferson Lab*

*Igor Strakovsky*  
イゴール ストラコフスキー

*The George Washington University*



- *KLF at Jefferson Lab.*
- *KLF experiment.*
- *Aims of KLF project.*
- *Impact to study Early Universe.*
- *Hyperon spectroscopy.*
- *Strange Meson spectroscopy.*
- *Where we are going.*
- *Summary.*
- *A bit of history.*

Supported by  DE-SC0016583




# KLF at Jefferson Laboratory





# Jefferson Lab *Continuous Electron Beam Accelerator Facility* in 2024



**1995 – 2012...**

Energy 0.4 – 6.0 GeV

- 200  $\mu\text{A}$ , Polarization 85%
- Simultaneous delivery 3 Halls – A, B, C

- 500+ PhDs completed
- On average 22 US Ph.Ds per year, roughly 25–30% of US Ph.Ds in nuclear physics
- 1530 users in FY16, ~1/3 international from 37 countries

**...2016 – ...**

Energy 0.4 – 12.0 GeV

- 150  $\mu\text{A}$ , Polarization 85%
- Simultaneous delivery 4 Halls
- FY18: First try simultaneous delivery to 4 Halls – A, B, C, D



Courtesy of Thia Keppel, 2017








## Strange Hadron Spectroscopy with Secondary $K_L$ Beam in Hall D

### Experimental Support:

Shankar Adhikari<sup>43</sup>, Moskov Amaryan (Contact Person, Spokesperson)<sup>13</sup>, Arshak Asaturyan<sup>1</sup>, Alexander Austregesilo<sup>49</sup>, Marouen Baalouch<sup>8</sup>, Mikhail Bashkanov (Spokesperson)<sup>63</sup>, Vitaly Baturin<sup>13</sup>, Vladimir Berdnikov<sup>11,35</sup>, Olga Cortes Becerra<sup>19</sup>, Timothy Black<sup>60</sup>, Werner Boeglin<sup>13</sup>, William Briscoe<sup>19</sup>, William Brooks<sup>54</sup>, Volker Burkert<sup>19</sup>, Eugene Chudakov<sup>19</sup>, Geraint Clash<sup>63</sup>, Philip Cole<sup>32</sup>, Volker Crede<sup>14</sup>, Donal Day<sup>61</sup>, Pavel Degtyarenko<sup>49</sup>, Alexandre Deur<sup>19</sup>, Sean Dobbs (Spokesperson)<sup>14</sup>, Gail Dodge<sup>13</sup>, Anatoly Dolgolenko<sup>26</sup>, Simon Eidelman<sup>6,41</sup>, Hovanes Egiyan (JLab Contact Person)<sup>49</sup>, Denis Epifanov<sup>6,41</sup>, Paul Eugenio<sup>14</sup>, Stuart Fegan<sup>63</sup>, Alessandra Filippi<sup>25</sup>, Sergey Furlotov<sup>19</sup>, Liping Gan<sup>60</sup>, Franco Garibaldi<sup>21</sup>, Ashot Gasparian<sup>39</sup>, Gagik Gavalian<sup>19</sup>, Derek Glazier<sup>18</sup>, Colin Gleason<sup>22</sup>, Vladimir Goryachev<sup>26</sup>, Lei Guo<sup>14</sup>, David Hamilton<sup>11</sup>, Avetik Hayrapetyan<sup>17</sup>, Garth Huber<sup>53</sup>, Andrew Hurley<sup>50</sup>, Charles Hyde<sup>13</sup>, Isabella Illari<sup>19</sup>, David Ireland<sup>18</sup>, Igal Jaegle<sup>49</sup>, Kyungseon Joo<sup>57</sup>, Vanik Kakoyan<sup>1</sup>, Grzegorz Kalicy<sup>11</sup>, Mahmoud Kamel<sup>13</sup>, Christopher Keith<sup>19</sup>, Chan Wook Kim<sup>19</sup>, Eberhard Klemp<sup>5</sup>, Geoffrey Krafft<sup>19</sup>, Sebastian Kuhn<sup>13</sup>, Sergey Kuleshov<sup>2</sup>, Alexander Laptev<sup>33</sup>, Ilya Larin<sup>26,39</sup>, David Lawrence<sup>19</sup>, Daniel Lersch<sup>14</sup>, Wenliang Li<sup>56</sup>, Kevin Luckas<sup>28</sup>, Valery Lyubovitskij<sup>50,51,52,54</sup>, David Mack<sup>49</sup>, Michael McCaughan<sup>49</sup>, Mark Manley<sup>30</sup>, Hrachya Marukyan<sup>1</sup>, Vladimir Matveev<sup>26</sup>, Mihai Mocanu<sup>63</sup>, Viktor Mokeev<sup>49</sup>, Curtis Meyer<sup>9</sup>, Bryan McKinnon<sup>18</sup>, Frank Nerling<sup>15,16</sup>, Matthew Nicol<sup>63</sup>, Gabriel Niculescu<sup>27</sup>, Alexander Ostrovidov<sup>14</sup>, Zisis Papandreou<sup>53</sup>, KiJun Park<sup>19</sup>, Eugene Pasyuk<sup>49</sup>, Peter Pauli<sup>18</sup>, Lubomir Pentchev<sup>19</sup>, William Phelps<sup>10</sup>, John Price<sup>7</sup>, Jörg Reinhold<sup>13</sup>, James Ritman (Spokesperson)<sup>28,68</sup>, Dimitri Romanov<sup>20</sup>, Carlos Salgado<sup>10</sup>, Todd Satogata<sup>49</sup>, Susan Schadmand<sup>28</sup>, Amy Schertz<sup>56</sup>, Axel Schmidt<sup>19</sup>, Daniel Sober<sup>11</sup>, Alexander Somov<sup>49</sup>, Sergei Somov<sup>35</sup>, Justin Stevens (Spokesperson)<sup>56</sup>, Igor Strakovsky (Spokesperson)<sup>19</sup>, Victor Tarasov<sup>26</sup>, Simon Taylor<sup>49</sup>, Annika Thiel<sup>5</sup>, Guido Maria Urciuoli<sup>24</sup>, Holly Szumila-Vance<sup>19</sup>, Daniel Watts<sup>63</sup>, Lawrence Weinstein<sup>13</sup>, Timothy Whitlatch<sup>49</sup>, Nilanga Wickramaarachchi<sup>13</sup>, Bogdan Wojtsekhowski<sup>19</sup>, Nicholas Zachariou<sup>63</sup>, Jonathan Zarling<sup>53</sup>, Jixie Zhang<sup>61</sup>

### Theoretical Support:

Alexey Anisovich<sup>5,44</sup>, Alexei Bazavov<sup>38</sup>, Rene Bellwied<sup>21</sup>, Veronique Bernard<sup>12</sup>, Gilberto Colangelo<sup>3</sup>, Aleš Cieplý<sup>16</sup>, Michael Döring<sup>19</sup>, Ali Eskanderian<sup>19</sup>, Jose Goity<sup>20,49</sup>, Helmut Haberzettl<sup>19</sup>, Mirza Hadžimehmedović<sup>55</sup>, Robert Jaffe<sup>36</sup>, Boris Kopeliovich<sup>54</sup>, Heinrich Leutwyler<sup>3</sup>, Maxim Mai<sup>19</sup>, Terry Mart<sup>65</sup>, Maxim Matveev<sup>44</sup>, Ulf-G. Meißner<sup>5,29</sup>, Colin Morningstar<sup>9</sup>, Bachir Moussallam<sup>12</sup>, Kanzo Nakayama<sup>58</sup>, Wolfgang Ochs<sup>37</sup>, Youngseok Oh<sup>31</sup>, Rifat Omerovic<sup>55</sup>, Hedim Osmanovic<sup>55</sup>, Eulogio Oset<sup>62</sup>, Antimo Palano<sup>64</sup>, Jose Peláez<sup>34</sup>, Alessandro Pilloni<sup>66,67</sup>, Maxim Polyakov<sup>18</sup>, David Richards<sup>49</sup>, Arkaitz Rodas<sup>19,56</sup>, Dan-Olof Riska<sup>12</sup>, Jacobo Ruiz de Elvira<sup>3</sup>, Hui-Young Ryu<sup>15</sup>, Elena Santopinto<sup>23</sup>, Andrey Sarantsev<sup>5,14</sup>, Jugoslav Stahov<sup>55</sup>, Alfred Švarc<sup>17</sup>, Adam Szczepaniak<sup>22,49</sup>, Ronald Workman<sup>19</sup>, Bing-Song Zou<sup>1</sup>



Extensive Theoretical Support

arXiv:2008.08215v2 [nucl-ex] 14 Sep 2020



Jefferson Lab PAC48 Report, 2020

**Summary:** The future  $K_L$  facility will add a new physics reach to JLab, and the PAC is looking forward to see the idea being materialized, in conjunction with the plans for Hall D as spelled out in the 2019 White Paper. The collaboration should now devote all its energy to turn this challenging project into an experimental facility and in parallel prepare for a successful data analysis.

e-Print: 2008.08215 [nucl-ex]


[https://wiki.jlab.org/klproject/index.php/Main\\_Page](https://wiki.jlab.org/klproject/index.php/Main_Page)



E12-12-19-001 *This Happens because of Strong Support & Dedicated Efforts of Collaboration*




# KLF Experiment







# Hall D Beam Line for




$$\frac{N(K_L) \text{ Jefferson Lab}}{N(K_L) \text{ SLAC}} = 10^3$$





12 GeV 5 μA  
Bunch spacing 128 ns

We will not use Pair Spectrometer




## Compact Photon Source




D. Day et al. Nucl Instrum Meth A 957, 163429 (2020)

## Kaon Production Target




## Kaon Flux Monitor



- Superior **CEBAF** electron beam will enable flux on order of  $10^4$   $K_f/sec$ , which exceeds flux of that previously attained @ **SLAC** by *three orders* of magnitude.

## Experimental Hall [GlueX Spectrometer, KFM, SC, Cryo Target]



## Tagger Hall [CPS]






## Collimator Cave [KPT]





# $K_L$ Momentum Determination & Beam Resolution

$$K_L^0 = \frac{1}{\sqrt{2}}(K^0 - \bar{K}^0)$$




Reconstruction of final-state particles

- Momentum measured with *TOF* between *SC* (surrounded *LH<sub>2</sub>/LD<sub>2</sub>*) & *RF* from *CEBAF*.
- *Mean lifetime* of  $K_L$  is 51.16 nsec ( $c\tau = 15.3$  m) whereas *Mean lifetime* of  $K^-$  is 12.38 nsec ( $c\tau = 3.7$  m).
- For this reason, it is much easier to perform measurements of  $K_L p$  scattering @ low beam energies compared with  $K^- p$  scattering.



# Electron Beam Parameters

- $E_e = 12$  GeV      $I = 5$   $\mu$ A
- Bunch spacing 64 vs 128 ns



- 128 ns confirmed feasible









# Aims of KLF Project



# We Can Do It, but Why?


- Why to use *kaon beam*? What is advantage compared to *electrons* or *photons* ?
- What is so special about *K-long* compared to *charged kaon* beam ? Complimentary to  .
- What is advantage of producing secondary kaon beam with *EM* probe, compared to *proton* beam ?
- How much  accelerator could make breakthrough compared to previous results @ **SLAC** ?
- Why to do this  experiment, what are we going to learn ?
- How will it affect our knowledge on *Hyperon Spectroscopy* ?
- What are we going to learn about *Strange Meson Spectroscopy* ?
- Is this  experiment about “*stamp collection*” or what ?

• There are many more *questions* - some constructive & some less so - answers to which shaped approved  experiment.










-  project has firmly to setup secondary  $K_L$  beamline @ **Jefferson Lab**, with *flux* of *three order of magnitude higher* than **SLAC** had, for scattering experiments on both *proton* & *neutron* (*first time !*) targets.
- **CEBAF** will remain *prime facility* for fixed target electron scattering @ luminosity *frontier*. *First hadronic facility* @ **Jefferson Lab**.

- We will determine differential cross sections & self-polarization of *hyperons* with **GlueX** detector to enable precise *PWA* in order to determine *all resonances* up to 2500 MeV in spectra of  $\Lambda^*$ ,  $\Sigma^*$ ,  $\Xi^*$ , &  $\Omega^*$ .  
To complete  $SU(3)_F$  multiplets, one needs no less than 48  $\Lambda^*$ , 38  $\Sigma^*$ , 61  $\Xi^*$ , & 31  $\Omega^*$ .

- We intend to do *strange meson spectroscopy* by studies of  $\pi$ - $K$  interaction to locate *pole* positions in  $I = 1/2$  &  $3/2$  channels.

-  has link to *ion-ion high energy* facilities such as  &  & will allow understand formation of our world in *several microseconds* after *Big Bang*. *Hyperons* are playing *leading* role to reproduce *Chemical Potential*.




Josiah Willard Gibbs





# *Impact to Study Early Universe*







 is *Home* of  
*Hot Big Bang Theory.*


- Omission of any “*missing hyperon states*” in *Standard Model* will negatively impact our understanding of *QCD freeze-out* in heavy-ion & hadron collisions, *hadron spectroscopy*, & *thermodynamics of Early Universe*.
- For that reason, advancing our understanding of formation of *baryons* from *quarks & gluons* requires new experiments to search for any *missing hyperon* resonances.



# Thermodynamics @ Freeze-Out

- In *thermodynamics*, *chemical potential* of *species* is *energy* that can be absorbed or released due to change of particle number of given species, *e.g.*, in chemical reaction or phase transition.
- *Chemical potential* of species in mixture is defined as rate of change of free energy of thermodynamic system with respect to change in number of atoms of species that are added to system.
- @ *chemical equilibrium* or in *phase equilibrium*, total sum of product of *chemical potentials* & stoichiometric coefficients is zero, as free energy is @ minimum.




Josiah Willard Gibbs

- Recent studies that compare *LQCD* calculations of *thermodynamic*, statistical *Hadron Resonance Gas* models, & ratios between measured yields of different hadron species in heavy ion collisions provide indirect evidence for presence of "missing" resonances in all these contexts.



Rolf Hagedorn



Chemical Potential

$$\left(\frac{\mu_S}{\mu_B}\right)_{LO} = -\frac{\chi_{11}^{BS}}{\chi_2^S} - \frac{\chi_{11}^{QS}}{\chi_2^S} \frac{\mu_Q}{\mu_B}$$

- + "Missing" Resonances (QM/LQCD calculations).

Contribution order:

- Hyperons
- Non-strange Baryons
- Mesons
- Light Nuclei

- Contribution from *observed Resonances*. PDG


- The *Early Universe* was *Strange*.



Courtesy of Claudia Ratti, YSTAR2016



# Hyperon Spectroscopy







*It is clear that we still need much more information about the existence and parameters of many baryon states, especially in the  $N=2$  mass region, before this question of non-minimal  $SU(6) \times O(3)$  super-multiplet can be settled.*

**Dick Dalitz, 1976**

*The first problem is the notion of a resonance is not well defined. The ideal case is a narrow resonance far away from the thresholds, superimposed on slowly varying background. It can be described by a **Breit-Wigner** formula and is characterized by a pole in the analytic continuation of the partial wave amplitude into the low half of energy plane.*



**Gerhard Höhler, 1987**



*Why  $N^*$ s are important – The first is that nucleons are the stuff of which our world is made. My second reason is that they are simplest system in which the quintessentially non-**Abelian** character of QCD is manifest. The third reason is that history has taught us that, while relatively simple, Baryons are sufficiently complex to reveal physics hidden from us in the mesons.*

**Nathan Isgur, 2000**




# Baryon Sector @ PDG2022

GW Contribution

R.L. Workman *et al*, Prog Theor Exp Phys 2022, 083C01 (2022)

$p$	$1/2^+$ ****	$\Delta(1232)$	$3/2^+$ ****	$\Sigma^+$	$1/2^+$ ****	$\Xi^0$	$1/2^+$ ****	$\Lambda_c^+$	$1/2^+$ ****
$n$	$1/2^+$ ****	$\Delta(1600)$	$3/2^+$ ***	$\Sigma^0$	$1/2^+$ ****	$\Xi^-$	$1/2^+$ ****	$\Lambda_c(2595)^+$	$1/2^-$ ***
$N(1440)$	$1/2^+$ ****	$\Delta(1620)$	$1/2^-$ ****	$\Sigma^-$	$1/2^+$ ****	$\Xi(1530)^0$	$3/2^+$ ****	$\Lambda_c(2625)^+$	$3/2^-$ ***
$N(1520)$	$3/2^-$ ****	$\Delta(1700)$	$3/2^-$ ****	$\Sigma(1385)$	$3/2^+$ ****	$\Xi(1620)^0$	*	$\Lambda_c(2765)^+$	*
$N(1535)$	$1/2^-$ ****	$\Delta(1750)$	$1/2^+$ *	$\Sigma(1480)$	*	$\Xi(1690)^0$	***	$\Lambda_c(2890)^+$	$5/2^+$ ***
$N(1650)$	$1/2^-$ ****	$\Delta(1900)$	$1/2^-$ **	$\Sigma(1560)$	**	$\Xi(1820)^0$	**	$\Lambda_c(2940)^+$	***
$N(1675)$	$5/2^-$ ****	$\Delta(1905)$	$5/2^+$ ****	$\Sigma(1580)$	$3/2^-$ *	$\Xi(1990)^0$	***	$\Sigma_c(2455)$	$1/2^+$ ****
$N(1690)$	$5/2^+$ ****	$\Delta(1910)$	$1/2^+$ ****	$\Sigma(1620)$	$1/2^-$ **	$\Xi(2030)^0$	$\geq 3/2^+$ ****	$\Sigma_c(2520)$	$3/2^+$ ****
$N(1695)$	*	$\Delta(1920)$	$3/2^-$ **	$\Sigma(1660)$	$1/2^+$ ***	$\Xi(2090)^0$	*	$\Sigma_c(2800)$	***
$N(1700)$	$3/2^-$ ***	$\Delta(1930)$	$1/2^-$ **	$\Sigma(1670)$	$3/2^-$ ****	$\Xi(2250)^0$	**	$\Xi_c^+$	$1/2^+$ ****
$N(1710)$	$1/2^+$ **	$\Delta(1940)$	$3/2^-$ **	$\Sigma(1690)$	**	$\Xi(2370)^0$	*	$\Xi_c^0$	$1/2^+$ ****
$N(1770)$	$3/2^+$ **	$\Delta(1950)$	$7/2^+$ **	$\Sigma(1750)$	**	$\Xi(2500)^0$	*	$\Xi_c^{*-}$	$1/2^+$ ****
$N(1830)$	$5/2^+$ **	$\Delta(2000)$	$5/2^+$ **	$\Sigma(1770)$	$1/2^+$ **	$\Omega(1670)^-$	$3/2^+$ **	$\Xi_c^0$	$1/2^+$ ****
$N(1880)$	$3/2^-$ **	$\Delta(2000)$	$1/2^-$ **	$\Sigma(1775)$	$1/2^-$ ****	$\Omega(2010)^-$	**	$\Xi_c(2645)$	$3/2^+$ ****
$N(1900)$	$7/2^-$ **	$\Delta(2200)$	$7/2^-$ **	$\Sigma(1840)$	$3/2^+$ **	$\Omega(2010)^-$	**	$\Xi_c(2790)$	$1/2^-$ ****
$N(1905)$	$1/2^-$ **	$\Delta(2300)$	$9/2^+$ **	$\Sigma(1880)$	$1/2^+$ **	$\Omega(2010)^-$	**	$\Xi_c(2815)$	$3/2^-$ ****
$N(1900)$	$1/2^-$ **	$\Delta(2350)$	$5/2^-$ *	$\Sigma(1910)$	$5/2^+$ ****	$\Omega(2010)^-$	**	$\Xi_c(2930)$	*
$N(1900)$	$7/2^-$ **	$\Delta(2390)$	$7/2^+$ *	$\Sigma(1910)$	$1/2^-$ **	$\Omega(2010)^-$	**	$\Xi_c(2980)$	***
$N(2000)$	$5/2^+$ **	$\Delta(2400)$	$9/2^-$ **	$\Sigma(2000)$	$1/2^-$ **	$\Omega(2010)^-$	**	$\Xi_c(3055)$	**
$N(2040)$	$3/2^+$ **	$\Delta(2420)$	$11/2^+$ ****	$\Sigma(2030)$	$7/2^+$ ****	$\Omega(2010)^-$	**	$\Xi_c(3080)$	***
$N(2060)$	$5/2^-$ **	$\Delta(2750)$	$13/2^-$ **	$\Sigma(2070)$	$5/2^+$ **	$\Omega(2010)^-$	**	$\Xi_c(3123)$	*
$N(2100)$	$1/2^+$ **	$\Delta(2950)$	$15/2^+$ **	$\Sigma(2080)$	$3/2^+$ **	$\Omega(2010)^-$	**	$\Omega_c^0$	$1/2^+$ ****
$N(2120)$	$3/2^-$ **			$\Sigma(2100)$	$7/2^-$ **	$\Omega(2010)^-$	**	$\Omega_c(2770)^0$	$3/2^+$ ****
$N(2190)$	$7/2^-$ ****	$\Lambda$	$1/2^+$ ****	$\Sigma(2250)$	***			$\Xi_c^+$	*
$N(2220)$	$9/2^+$ ****	$\Lambda(1405)$	$1/2^-$ ****	$\Sigma(2455)$	**			$\Xi_c^0$	*
$N(2250)$	$9/2^-$ ****	$\Lambda(1520)$	$3/2^-$ ****	$\Sigma(2620)$	**			$\Lambda_b^0$	$1/2^+$ ***
$N(2600)$	$11/2^-$ ****	$\Lambda(1600)$	$1/2^+$ ***	$\Sigma(3000)$	*			$\Sigma_b^+$	$1/2^+$ ****
$N(2700)$	$13/2^+$ **	$\Lambda(1670)$	$1/2^-$ ****	$\Sigma(3170)$	*			$\Sigma_b^0$	$3/2^+$ ****
		$\Lambda(1690)$	$3/2^-$ ****					$\Xi_b^+$	$1/2^+$ ****
		$\Lambda(1800)$	$1/2^-$ ****					$\Xi_b^0$	$1/2^+$ ****
		$\Lambda(1810)$	$1/2^-$ ****					$\Xi_b^+$	$1/2^+$ ****
		$\Lambda(1820)$	$5/2^+$ ****					$\Xi_b^0$	$1/2^+$ ****
		$\Lambda(1830)$	$5/2^-$ ****						
		$\Lambda(1890)$	$3/2^+$ ****						
		$\Lambda(2000)$	$1/2^-$ ****						
		$\Lambda(2000)$	$7/2^+$ ****						
		$\Lambda(2100)$	$7/2^-$ ****						
		$\Lambda(2110)$	$5/2^+$ ****						
		$\Lambda(2325)$	$3/2^-$ **						
		$\Lambda(2350)$	$9/2^+$ **						
		$\Lambda(2585)$	**						



- PDG2022 has 133 Baryon Resonances (69 of them are 4\* & 3\*).
- In case of SU(6) x O(3), 434 states would be present if all revealed multiplets were fleshed out (three 70 & four 56).
- LQCD results are similar.

R. Koniuk & N. Isgur, Phys Rev Lett 44, 845 (1980)

• First hyperon was discovered in 1950.

• Pole position in complex energy plane for hyperons has been made only in 2010.




V.D. Hopper & S. Biswas, Phys Rev 80, 1099 (1950)


Y. Qung *et al*, Phys Lett B 694, 123 (2010)



# Baryon Multiplets of Eight-fold Way




- Three light quarks can be arranged in 6 baryonic families,  $N^*$ ,  $\Delta^*$ ,  $\Lambda^*$ ,  $\Sigma^*$ ,  $\Xi^*$ , &  $\Omega^*$ .
- Number of members in family that can exist is not arbitrary.
- If  $SU(3)_F$  symmetry of QCD is controlling, then:



← Spin 1/2 baryon octet:  $N^*$ ,  $\Lambda^*$ ,  $\Sigma^*$ ,  $\Xi^*$

Spin 3/2 baryon decuplet:  $\Delta^*$ ,  $\Sigma^*$ ,  $\Xi^*$ ,  $\Omega^*$  →



Resonance	LQCD	Observed
$N^*$	62	36
$\Delta^*$	38	29
$\Lambda^*$	71	23
$\Sigma^*$	66	28
$\Xi^*$	73	12
$\Omega^*$	36	5



R. G. Edwards et al, Phys Rev D **87**, 054506 (2013)

- Seriousness of “missing-states” problem is obvious from these numbers.
- One needs to complete  $SU(3)_F$  multiplets.



R. Koniuk & N. Isgur, Phys Rev Lett **44**, 845 (1980)




B.M.K. Nefkens,  $\pi N$  Newsletter, **14**, 150 (1997)






# Road Map to Baryon Spectroscopy


• That is not hunting for bumps



Data




Amplitudes



PWA

PWA





Chen et al. [Particle Data Group], Chin. Phys. C, 40, 10301 (2016) and 2017 update

$\Delta(1232) \ 3/2^+$   $J^P = \frac{3}{2}^+$  Status: \*\*\*

Other and absolute values are listed and referenced in the 2014 edition, Chinese Physics C38 070001 (2014)

$\Delta(1232)$  POLE POSITIONS

REAL PART, MIXED CHARGES	DOCUMENT ID	TECH.	COMMENT
1209 $m = 1$	1211	1211	1211
1211 $m = 1$	1211	1211	1211
1215.5-1.0	1211	1211	1211
1216	1211	1211	1211
1219	1211	1211	1211
1219 $m = 1$	1211	1211	1211





- Limited number of  $K_L$  induced measurements (1961 – 1982)  
 $2426 d\sigma/d\Omega$ ,  $348 \sigma^{\text{tot}}$ , &  $115 P$  observables do not allow today to *feel comfortable* with *Hyperon Spectroscopy* results.


$W = 1.45 - 5.05$  GeV

- Limited number of  $K_L$  observables in *hyperon spectroscopy* @ present poorly constrain phenomenological analyses.

- Overall systematics* of previous experiments varies between **15%** & **35%**.  
*Energy binning* is much broader than hyperon widths.


- There were **no measurements using polarized target**. It means that there are no *double polarized* observables which are critical for *complete experiment* program.

- We are not aware of any data on *neutron* target.




Target  $\rightarrow$  *Proton*


Elastic & Charge-Exchange




Two-body with  $S = -1$




Two-body with  $S = -2$








Three-body with  $S = -2$





Three-body with  $S = -3$



*Neutron* [first measurements]



- To search for “missing” hyperons, we need measurements of production reactions:







# Samples of PWA Results for Current DB

H. Zhang et al Phys Rev C 88, 035204 (2013)

H. Zhang et al Phys Rev C 88, 035205 (2013)





• PWA (  &  ) predictions at lower & higher energies have poorer agreement for  $S \neq 0$  data than for  $S = 0$  data.


• Polarized measurements are *tolerable* for any PWA solutions.



# Impact Proposed Data using PWA








Quasi-data using *GlueX* detector properties  



100 days  
 20 days

• At least **100** days needed to get *precise* solution.


Resonance	 20 days: M, $\Gamma$	 100 days: M, $\Gamma$	 PDG: M, $\Gamma$	 had <sub>spec</sub> M
$\Sigma(1920)5/2^-$	1977 $\pm$ 21 $\pm$ 25 327 $\pm$ 25 $\pm$ 25	1923 $\pm$ 10 $\pm$ 10 321 $\pm$ 10 $\pm$ 10		2027 2487 2659 2781
$\Sigma(2030)7/2^+$	1981 $\pm$ 30 $\pm$ 30 350 $\pm$ 80	1930 $\pm$ 20 $\pm$ 30 400 $\pm$ 40	2030 $\pm$ 10 180 $\pm$ 30	2686 2709 2793 2806



# Theory for "Neutron" Target Measurements



- There are 6 different models.
- Pole positions of  $\Lambda(1405)$  in chiral unitary approaches.
- Each symbol represents position of 1<sup>st</sup> (black) & 2<sup>nd</sup> (red) pole in each model.



•  $P = 300$  MeV/c




Courtesy of Maxim Mai, 2019



# Summary of Hyperon Spectroscopy


Money Plot



• We showed that  $K_L$  sensitivity with **100 days** of running will allow to discovery many *hyperons* with good precision.

• *Why should it be done with KL beam ?*

This is only realizable way to observe s-channel resonances having *all momenta* of  $K_L$  @ once ("tagged" kaons).



• *Why should it be done @ Jefferson Lab ?*

Because nowhere else in existing facilities this can be done.

• *Why should we care that there are dozens of missing states ?*

...The new capabilities of the **12-GeV** era facilitate a detailed study of baryons containing two and three strange quarks. Knowledge of the spectrum of these states will further enhance our understanding of the manifestation of **QCD** in the three-quark arena.

**2015 Long Range Plan for Nuclear Science**





# Narrow Pentaquarks from $\Lambda_6 \rightarrow J/\psi p \bar{K}^-$

- QCD gives rise to *hadron spectrum*.

Volume 8, number 3      PHYSICS LETTERS      1 February 1964

A SCHEMATIC MODEL OF BARYONS AND MESONS      CI=4087

M. GELL-MANN

If we assume that the strong interactions of baryons and mesons are correctly described in terms of the broken "eightfold way"      Baryons can now be constructed from quarks by using the combinations  $(qqq)$ ,  $(qqqqq)$ , etc., while mesons are made out of  $(q\bar{q})$ ,  $(qq\bar{q}\bar{q})$ , etc.

- Many  $\bar{q}q$  &  $qqq$  states have been observed.


PDG 220 & 100.

- $\bar{q}q\bar{q}q$ ,  $qqq\bar{q}q$ ... are *not forbidden* or we do not know it yet.

- LHCb claims evidence for *four hidden-charm  $qqq\bar{q}q$  states* near *open-charm* decay thresholds for  $\Sigma_c^+ \bar{D}^0$  &  $\Sigma_c^+ \bar{D}^{*0}$  in  $\Lambda_6 \rightarrow J/\psi p \bar{K}^-$  decays.

## Bump hunting:

- no quantum numbers
- no pole positions



R. Aaij *et al*, Phys Rev Lett 122, 222001 (2019)


CI=666



State	M (MeV)	$\Gamma[P_c \rightarrow J/\psi + p]$ (MeV)	Significance
$P_c(4312)^+$	$4311.9 \pm 0.7^{+6.8}_{-0.6}$	$9.8 \pm 2.7^{+3.7}_{-4.5}$	$7.3 \sigma$
$P_c(4337)^+$	$4337^{+7+2}_{-4-2}$	$29^{+26+13}_{-12-14}$	$3.1 - 3.7 \sigma$
$P_c(4440)^+$	$4440.3 \pm 1.3^{+4.1}_{-4.7}$	$20.6 \pm 4.9^{+8.7}_{-10.1}$	$5.4 \sigma$
$P_c(4457)^+$	$4457.3 \pm 0.6^{+4.1}_{-1.7}$	$6.4 \pm 2.0^{+5.7}_{-1.9}$	$5.4 \sigma$



# How Bump Hunting works in 2019 GLUEX citations experiment data?




A. Ali *et al*, Phys Rev Lett **123**, 072001 (2019)



CI=175

2016–2017 data:  $469 \pm 22 \gamma p \rightarrow J/\psi p \rightarrow e^+e^-p$  &  $68 \text{ pb}^{-1}$



M.J. Amarian, Few-Body Syst Suppl. **11**, 359 (1999)

A. Ali *et al*, Phys Rev Lett **123**, 072001 (2019)

U. Camerini *et al*, Phys Rev Lett **35**, 483 (1975)


B. Gittelman *et al*, Phys Rev Lett **35**, 1616 (1975)

D. Kharzeev, H. Satz, A. Syamtomov, & G. Zinovjev, Nucl Phys A **661**, 568 (1999)

JPAC A.N. Hiller Blin *et al*, Phys Rev D **94**, 034002 (2016)



S. Brodsky, E. Chudakov, P. Hoyer, & J.M. Laget, Phys Lett B **498**, 23 (2001)

• 3g works better than 2g @ threshold




• GLUEX sees *no evidence* for LHCb  $P_c$ s  
Upper limits @ 90% CL

State	Upper Limit
$P_c(4312)$	4.6 %
$P_c(4440)$	2.3 %
$P_c(4457)$	3.8 %



Assuming  $J^P = 3/2^-$






# Alternative Solution for **GLUEX** Data

IIS, W.J. Briscoe, E. Chudakov, I. Larin, L. Pentchev, A. Schmidt, R.L. Workman,  
 Phys Rev C **108**, 015202 (2023)

- We suggested to apply *rearrangement interference* for revealing *faint* resonance signals (*amplification* by *interference* with *strong* background signal).
- Relative phase  $\alpha$  leads to *constructive* (*bump*) or *destructive* (*dip*) *interference* for particular **PW**.

$$f = b + R \cdot \exp(2i\alpha)$$



**Resonance:**  $\chi^2/ndf=11.99/12=1.00$   
 $M = 4235 \pm 8$  MeV  
 $\Gamma = 35.4 \pm 8.2$  MeV *Resolution ~6 MeV*  
 $X = 0.023 \pm 0.005$   
 $\alpha = 40.8 \pm 5.7$  deg

**Background:**  
 $A = 0.00251 \pm 0.00046$  nb GeV/c  
 $B = 0.00688 \pm 0.00083$  nb/GeV/c

**No Resonance:**  $\chi^2/ndf=19.74/16=1.23$   
 $A = 0.00183 \pm 0.00040$  nb GeV/c  
 $B = 0.00766 \pm 0.00077$  nb/GeV/c

- *Dip* position does not correspond to *real mass* of  $P_c(4312)^+$ .
- It may depend on reaction *mechanism* (including *cusps* (*open charm*)) & background choices.

- If “*bump*” is imposed on **GLUEX** data “*by hand*” (consider **7th - 9th** energy values up from threshold), qualitative description of data up to  $W = 4.35$  GeV is possible, but with higher  $\chi^2$ , if our fit form is used.


- Obtained mass in our analysis is almost **77 MeV** below **LHCb** determination, but it cannot exclude that this is  $P_c(4312)^+$ .




Anyone can ask *Big Questions*, but it is not easy to ask questions that would suggest new pathways leading to real progress of our understanding.

Courtesy of Gerard 't Hooft, 2022

## 10 & $\bar{10}$ - P wave Multiplets



J.J. de Swart, Rev Mod Phys 35, 916 (1963)



$K_L p \rightarrow K^+ K^+ K^- E^{--}$

$K_L p \rightarrow K^+ n$

$K_L p \rightarrow K_S E^+$

Not N(1710)  
R.A. Arndt et al.  
Phys Rev C 69, 035208 (2004)



• In addition to ordinary hyperon states made of  $qqq$ , experiment may observe exotic  $\Theta^+$  pentaquark lying in apex of  $\bar{10}$  it will also be sensitive to observe exotic  $E^+$  lying in *right corner* of  $\bar{10}$  in reaction  $K_L p \rightarrow K_S E^+$  & to observe another exotic state  $E^-$  lying in *left corner*  $\bar{10}$  in reaction  $K_L p \rightarrow K^+ K^+ K^- E^{--}$



D. Diakonov, V. Petrov, & M.V. Polyakov, Z. Phys. A 359, 305 (1997)

CI=987





- Originally PWA arose as technology to determine amplitude of reaction via fitting scattering data.

⇒ That is *non-trivial mathematical problem* – looking for solution of **ill-posed** problem following to Hadamard & Tikhonov.

[number of equations less than number of unknown quantities]

⇒ There are two main technologies to look for solution:

(i) *least-squares minimization* of functions which are linear in unknown parameters,  $\chi^2$  &

(ii) *likelihood measures goodness* of fit of statistical model.

[Minimizing  $\chi^2$  is equivalent to maximizing (log) likelihood just case not small statistics]

⇒ Model *independent* treatment or data *driven* treatment.



Roger Cotes



Sir Ronald Aylmer Fisher

- Resonances appeared as by-product

[bound states objects with definite quantum numbers, mass, lifetime, & so on].

- Standard PWA

⇒ Reveals only wide Resonances, but not too wide ( $\Gamma_R < 500$  MeV) & possessing not too small BR (BR > 4%).

⇒ Tends (by construction) to miss narrow Resonances with  $\Gamma_R < 20$  MeV.




# Standard $KN$ PWA

J.S. Hyslop, R.A. Arndt, L.D. Roper, & R.L. Workman, Phys Rev D **46**, 961 (1992)



CI=163




$T = 0$  [20] 1100 MeV



**Pole Positions:**

I	Ampl	ReW (MeV)	-ImW (MeV)
0	$P_{01}$	1831	95
	$D_{03}$	1788	170
1	$P_{13}$	1811	118
	$D_{15}$	2074	253

**All Res in standard PWA are too heavier & too broader than  $\Theta^+$ .**



$K^+n \rightarrow K^+n$	- 98
$K^+n \rightarrow K^0p$	- 6
$K^+d \rightarrow K^0pp$	- 77
$K^+d \rightarrow K^+np$	- 43

*Modified  $KN$  PWA for  $\Theta(1540)$*   
 R.A. Arndt, IIS, & R.L. Workman, Phys Rev C **68**, 042201(R) (2003)





- Because PWA (by construction) tends to miss narrow Res with  $\Gamma_R < 20$  MeV
- We assume existence of Res & refit over whole DB

- Insertion of narrow Res in PWA for

Elastic case:  $e^{2i\delta} \Rightarrow e^{2i\delta}_R e^{2i\delta}_B$

$$e^{2i\delta}_R = (M_R - W + i\Gamma_R/2) / (M_R - W - i\Gamma_R/2)$$

Inelastic case:  $\eta e^{2i\delta} \Rightarrow \langle a|S|a \rangle = r_a A(W) e^{2i\delta}_R + (1 - r_a) B(W)$

$$r_a = BR(R \rightarrow a) \quad |A(M_R)| = 1 \quad \Sigma r_a = 1$$

$$\eta \leq 1 \Rightarrow r_a |A(W)| + (1 - r_a) |B(W)| \leq 1$$





## Refitting

- **Worse description**
    - $\Rightarrow$  Res with corresponding  $M_R$  &  $\Gamma_R$  is not supported
  - **Better description**
    - $\Rightarrow$  Res may exist
    - $\Rightarrow$  Effect can be due to various corrections (eg, thresholds)
    - $\Rightarrow$  Both possibilities can contribute
- Some additional checks are necessary
- True Res should provide effect only in particular PW
  - While Non-Res source may show similar effects in various PWs





# $\Delta\chi^2$ due to Insertion of Res into Different Waves

R.A. Arndt, IIS, & R.L. Workman, Phys Rev C **68**, 042201(R) (2003)



- Res contributes  $\sim \Gamma_{el} / (M_R - W)$  @  $|M_R - W| \gg \Gamma_R$
- For  $M_R \sim 1545$  MeV,  $\Gamma_R < 0.5$  MeV



**For  $I = 0$ :**


- Only one partial wave ( $P_{01}$ ) admits effect near 1545 MeV: resonance,  $\Gamma < 0.5$  MeV
- Other partial waves ( $S_{01}$  &  $P_{03}$ ) may have the effect only by accompanied by other corrections





# What One Can Expect for $K_L p \rightarrow K^+ n$ from ?

M. Amarian, S. Hirama, D. Jido, & IIS, e-Print: 2401.05887 [hep-ex]



V.V. Barmin *et al* Phys Rev C **89**, 045204 (2014):  $M = 1538 \pm 2$  MeV  $\rightarrow p_{KL} = 440$  MeV/c  $\rightarrow k = 0.268$  GeV/c &  $\Gamma = 0.34 \pm 0.10$  MeV

$K^+ X e \rightarrow K^0 p X e'$

Ghil-Seok Yang & Hyun-Chul Kim, PTEP **2013**, 013D01 (2013) ChSA:  $\Gamma = 0.5 \pm 0.1$  MeV

R.A. Arndt, IIS, & R.L. Workman, Phys Rev C **68**, 042201(R) (2003): Modified KN PWA gave:  $\Gamma < 0.5$  MeV @  $M \sim 1545$  MeV

Assuming  $\theta^+$  width  $\Gamma = 0.4$  MeV



R.N. Cahn & G.H. Trilling, Phys Rev D **69**, 011501(R) (2004)

$$\sigma_0 = \frac{2J+1}{(2s_{KL}+1)(2s_p+1)} \frac{4\pi}{k^2} = 68 \text{ mb}$$





Y. Iizawa, D. Jido, & S. Huebsch, [arXiv:2308.09397 [hep-ph]]: Assuming bkgd Xsec @ 440 MeV/c

$\sigma_0 = 3$  mb



e-Print: 2008.08215 [nucl-ex]


@ 1 MeV/c bin & flux =  $0.1$  (MeV/c s) $^{-1}$ .  
It means - we will have  $\sim 10^6$   $K_L$ /100 days.



$$\sigma_p/p = 6 \times 10^{-3}$$

10,000

$$N_{peak} = \frac{\Gamma(\theta^+) \pi \sigma_0 N_{bkgd} B_i B_f}{2\sigma_{bkgd} \Delta m_0} = 18,000 \text{ events}$$








18k events in peak in 100 days.  
Taking into account  
- acceptance &  
- efficiency of kaon registration,  
One can expect 100 events per day.


$$R = S/B = 4/1$$






We propose to search for  $\Theta^+$  in  $K^+ d \rightarrow K^0 pp$  reaction at  $p_{K^+}=0.5$  GeV/c at . A large acceptance Hyperon Spectrometer, which consists mainly of a time projection chamber and a 1-T superconducting magnet, will exclusively measure the decay products of  $\Theta^+$ , such that  $\Theta^+ \rightarrow K^0 p$ , followed by  $K^0 \rightarrow \pi^+ \pi^-$ , with a mass resolution of  $1$  MeV at  $M_{\Theta}$ . We investigated the feasibility of the proposed experiment using a Monte Carlo simulation. As a result, we expect to collect five orders of magnitude  $\Theta^+$  events, assuming a cross section of  $300 \mu\text{b}$  in 15-day beam time at .



■ @ 0.434 GeV/c C.J.S. Damerell et al, Nucl Phys B **94**, 374 (1975) 

● @ 0.470 GeV/c R.G. Glasser et al., Phys. Rev. D **15**, 1200 (1977) 

● Expectation is 137,000  $\Theta^+$  events



# *If $\Theta^+$ does not Survive, 'Damned' Questions Revive:*


- *Why are there no strongly bound exotic states..., like those of two quarks and two antiquarks or four quarks and one antiquark?*

H. Lipkin, Phys Lett **45B**, 267 (1973)




- *...either these states will be found by experimentalists, or our confined, quark-gluon theory of hadrons is as yet lacking in some fundamental, dynamical ingredient which will forbid the existence of these states or elevate them to much higher masses.*

R. Jaffe & K. Johnson, Phys Lett **60B**, 201 (1976)



# Strange Meson Spectroscopy








*The di-quark or meson-baryon puzzle: Why is the quark-quark interaction just enough weaker than the quark-anti-quark interaction so that di-quarks near the meson mass are not observed, but three-quark systems have masses comparable to those of mesons?*

**Harry Lipkin, 1973**

*For the region below 1 GeV, the debate centers on whether the phenomena are truly resonant or driven by attractive  $t$ -channel exchanges, and if the former, whether they are molecules or  $\bar{q}q\bar{q}q$ .*

**Frank Close, 2007**



*QCD predicts there should be a far richer spectrum, with states made predominantly of glue, we call glueballs, tetra-quark states made of two quarks and two anti-quarks... For almost forty years we have been searching for these additional states. Indeed, we may well have observed some of these, but there is little certainty of what has been found.*

**Michael Pennington, 2015**


*A simple picture for both mesons and baryons is inconsistent with any version of relativistic field theory, where one can not exclude presence of an arbitrary number of virtual  $\bar{q}q$  pairs and/or gluons. Therefore, adequate description of any hadron should use a Fock column, where lines correspond to particular configurations (but with the same "global" quantum numbers, like  $I, J, P, C$ , and so on).*

**Yakov Azimov, 2015**



# Scalar Meson Nonet

SU(3):  $J^{PC} = 0^{++}$  Nonet




- **Four** states called  $\kappa(700)$ .
- **still need further confirmation.**
- allows determination of all **four** states.

$$M = a_0 + a_1 Y + a_2 \left[ I(I+1) - \frac{1}{4} Y^2 \right]$$

$I$  – isospin,  $Y$  – strangeness (hypercharge),  $a_i$  – free prmts.

• This phenomenological formula works with accuracy of **5%**.



• **Mixing** be able to **shift** some masses for *Gell-Mann-Okubo mass* formula.



# Scalars vs Vectors or Eyewitness of 4q Exotics?



R.J. Jaffe, Phys Rev D **15**, 267 (1977)

arXiv: 0001123 [hep-ph]

Prog Theor Phys Suppl **168**, 127 (2007)

*Inverted mass hierarchy tetraquarks*

**Scalar Mesons**



- Very different mass *hierarchy*.
- Possibly suggesting *4q tetraquark*.
- Structure of *scalar* mesons.






- Certainly, there is no clear distinction between *4q* & "meson molecule" categories.




"I like the conclusion that the  $a_0$  is a multi quark state." Courtesy of Bob Jaffe, 2022



# Proposed Measurements for $K\pi$ Scattering



Unmeasured



region of  $K(800)$


SLAC Lower limit






# Summary of $K\pi$ Spectroscopy

Money Plot



-  will have very significant *impact* on our knowledge on  $K\pi$  scattering amplitudes.
- It will certainly improve still conflictive determination of *heavy  $K^*$ 's parameters*.
- It will help to settle tension between phenomenological determination of *scattering lengths* from data vs *ChPT & LQCD*.
- For  $K^*(700)$ , it will reduce:
  - *uncertainties* in *mass* by factor of *two* &
  - *uncertainties* in *width* by factor of *five*.
- It will help to clarify debated of its *existence*, &, therefore, *long standing problem* of existence of *scalar meson nonet*.

 100 days

• **Roy-Steiner** dispersion approach  
 $M - i\Gamma/2 = (648 \pm 4) - i(280 \pm 8) \text{ MeV}$



J.R. Pelaez *et al* Phys Rev D **93**, 074025 (2016)



# Where We are Going



# Beam Time Approved by PAC48

- Expected cornucopia of differential cross sections of different reactions with  $LH_2$  & below  $W = 2.5$  GeV for 100 days of beam time:

For  $d\sigma/d\Omega$

Reaction	Statistics (events)
$K_{LP} \rightarrow K_{SP}$	2.7M
$K_{LP} \rightarrow \pi^+\Lambda$	7M
$K_{LP} \rightarrow K^+\Xi^0$	2M
$K_{LP} \rightarrow K^+n$	60M
$K_{LP} \rightarrow K^-\pi^+p$	7M


For  $p$ , statistics is 0.2M

- There are no data on ``neutron'' targets &, for this reason, it is hard to make realistic estimate of statistics for  $K_L n$  reactions.  
If we assume similar statistics as on ``proton'' target, full program will be completed after running 100 days with  $LH_2$  & 100 days with  $LD_2$  cryo targets.
- Expected systematics for  $d\sigma/d\Omega$  is 10% or less.



# Jefferson Lab *Hall D Tentative Schedule*

*Safety Pauses Shifted Upcoming Schedule Out By Two Months And Is Not Yet Shown On This Figure.*



Courtesy of Doug Higinbotham, Sept 2023





# SUMMARY

Experiment was approved in **08/20**.  
ERR-I was approved in **08/23**,  
ERR-II may come in **06/24**.

- Our goal is
  - To setup *KL Facility* @ 
  - To measure & bring *new physics*.

-   would advance *Hyperon Spectroscopy* & study of *strangeness* in nuclear & hadronic physics.

*We may have cornucopia of many missing/new strange states.*

To complete  $SU(3)_F$  multiplets, one needs no less than **48  $\Lambda^*$ , 38  $\Sigma^*$ , 61  $\Xi^*$ , & 31  $\Omega^*$**



- Discovering of “*missing*” *hyperon states* would assist in advance our understanding of formation of *baryons* from *quarks* & *gluons* **microseconds** (!) after *Big Bang*.

Our expectation is to get **1** missed/new *hyperon* per **1** day.

- In *Strange Meson Spectroscopy*, *PWA* will allow to determine excited  $K^*$  states including *scalar  $K^*(700)$*  states.



桜が近づいてきました



ありがとうございました。

質問を歓迎します





**BACKUP**



# A Bit of History





PHYSICAL REVIEW

VOLUME 138, NUMBER 5B

7 JUNE 1965

First paper on subject

Photoproduction of Neutral K Mesons\*

CP-violation (1964)  
Hot topic!

S. D. DRELL AND M. JACOB†

Stanford Linear Accelerator Center, Stanford University, Stanford, California

(Received 6 January 1965)



Photoproduction of a neutral  $K$ -meson beam at high energies from hydrogen is computed in terms of a  $K^*$  vector-meson exchange mechanism corrected for final-state interactions. The results are very encouraging for the intensity of high-energy  $K_2$  beams at high-energy electron accelerators. A typical magnitude is  $20 \mu\text{b}/\text{sr}$  for a lower limit of the  $K^0$  photoproduction differential cross section, at a laboratory peak angle of  $2^\circ$ , for 15-BeV incident photons.




FIG. 1.  $K^*$  exchange in photoproduction.




FIG. 3. Center-of-mass differential cross section at 10 BeV. Curve (1) gives the Born approximation. Curve (2) is obtained after subtraction of the  $j=\frac{1}{2}$  partial wave. Curves (3) and (4) are respectively obtained after the  $j=\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \frac{7}{2}$ , and all partial waves have been corrected for absorption in final state. The results shown as directly obtained from  $600$  data by the computer.

Our motivation in carrying out this calculation is to emphasize the strong suggestion that an intense “healthy”  $K_2$  beam will emerge from high-energy electron accelerators (SLAC in particular) and will be available for detailed experimental studies.

Courtesy of Mike Albrow, KL2016

3/16/2024

Seminar, Tokyo Tech, Tokyo, February 2024

The possibility that useful  $K_L$  beam could be made @ electron synchrotron by photoproduction was being considered, & 1965 prediction for SLAC by Drell & Jacob was optimistic.



8.B.5 Nuclear Physics B23 (1970) 509-524. North-Holland Publishing Company  
8.B.6

## PHOTOPRODUCTION OF $K^0$ MESONS FROM PROTONS AND FROM COMPLEX NUCLEI

M. G. ALBROW<sup>†</sup>, D. ASTON, D. P. BARBER, L. BIRD<sup>††</sup>,  
R. J. ELLISON, C. HALLIWELL, A. E. HARCKHAM<sup>†††</sup>,  
F. K. LOEBINGER, P. G. MURPHY, J. WALTERS<sup>††</sup> and A. J. WYNROE  
*Schuster Laboratories, The University of Manchester,  
Manchester M13 9PL*



R. F. TEMPLEMAN  
*Daresbury Nuclear Physics Laboratory, Daresbury,  
Near Warrington, Lancs.*



Received 16 July 1970

Study photoproduction as means of making clean  $K^0$  beams & their decays & later, interactions.

**From: Mike Albrow** Aug 29, 2020  
**To: Igor Strakovsky**  
*Dear Igor, That is excellent news, thank you for letting me know. In one of those strange coincidences, my professor at Manchester who had the idea for our  $K^0$  photoproduction experiments and led the program, Paul Murphy (Manchester Univ.) died on Wednesday Aug 26. He was 89. I had told him about your plans, he was still interested. He would have been happy to know that 50 years later you are benefitting from his idea.  
Best, Mike (I am doing well, thank you)  
PS: If your proposal was accepted on Aug 26th let me know, it would be strange synchronicity!*

VOLUME 22, NUMBER 18

PHYSICAL REVIEW LETTERS

5 MAY 1969

## PRODUCTION OF $K_L^0$ MESONS AND NEUTRONS BY 10- AND 16-GeV ELECTRONS ON BERYLLIUM\*



A. D. Brody, W. B. Johnson, D. W. G. S. Leith, G. Loew, J. S. Loos, G. Luste, R. Miller, K. Moriyasu, B. C. Shen, W. M. Smart, and R. Yamartino

SLAC

Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305  
(Received 13 March 1969)

Systematics of particle-anti-particle processes through intrinsic property of K-longs.



# Kinematics for $K_{LP} \rightarrow K^+ n$ & $K_{LP} \rightarrow p K_S$

