

Nuclear gluons with charm at EIC

JLab FY16 LDRD Project LD1601

Collaborators: E. Chudakov, D. Higinbotham (10%), S. Furletov (20%/10%), C. Weiss (20%)
Ch. Hyde (4 weeks/year), M. Stratmann (2 weeks), M. Strikman (2 weeks)
Dien Nguyen (graduate student UVA & JLab), second student (2016)


Project period: 2 years

Aim: Demonstrate feasibility of direct probes of nuclear gluons & quarks with EIC:
Gluons at $x \gtrsim 0.1$ with heavy quarks — open charm, beauty;
Quark flavor decomposition with semi-inclusive DIS.


Tasks

- Adapt simulation tools for heavy quark production in ep (HVQDIS, F_{2c} code) to eA at EIC, with schematic modeling of charm detection/reconstruction, simple \rightarrow more detailed
- Adapt MC tool for ep semi-inclusive DIS to eA at EIC
- Simulate nuclear ratio measurements of open charm production and semi-inclusive DIS at EIC
- Assess systematic and theoretical accuracy of nuclear ratio measurements at EIC and quantify physics impact


Nuclear modification of partonic structure



$$\begin{array}{c} \updownarrow \\ \leftarrow \circ \rightarrow \\ \updownarrow \\ \dots \end{array} = |N\rangle + \sum |\text{non-}N\rangle$$



quark exchange



meson exchange

- Seen in inclusive DIS
JLab 6 & 12 GeV: EMC effect
- Gluonic EMC effect at $x > 0.3$?
Non-nucleonic DOF in nucleus
- Quark and gluon enhancement at $x \sim 0.1$ — antishadowing?

QCD structure of NN interaction


Flavor structure of quark enhancement?

Gluon enhancement?

Strong gluon shadowing observed at $x < 0.01$ suggests compensating antishadowing at $x \sim 0.1$

LHC ALICE J/ψ in ultraperipheral AA

- Need direct probes!



- Heavy quark production in DIS

Calculated in QCD at LO, NLO; estimates beyond NLO
 Laenen, Riemersma, Smith Van Neerven, Harris 93+. Kawamura et al. 12

Theoretical uncertainties studied: Scheme dep, quark mass
 Alekhin, Moch et al. 93+

Probes gluons at $x' > \frac{4M_Q^2 + Q^2}{W^2}$ ($W^2 \gg Q^2$)

Good sensitivity to gluons even at $x' \gtrsim 0.1$

- Experimental identification

Single D , B mesons through distinctive leptonic or hadronic decays, e.g. D^*

Vertex detection through track reconstruction or vertex detector

Double detection, e.g. $\Lambda_c + D$?

- Observables for analysis

Differential cross section $d^4\sigma/dQ^2 d\eta d^2p_T$

Inclusive charm structure function F_{2c}

- Experimental results

ep HERA H1 & ZEUS: extensive measurements, various identification methods, detailed comparison with QCD calculations

Latest analysis: Aaron et al. 2011, Abramowicz et al. 2014

μN COMPASS: polarized target

Adolph et al. 2012/13

Charm production in e^+e^- , photon-hadron, hadron-hadron

- Simulation tools

HVQDIS code: Calculates differential D -meson production cross section using NLO QCD approximation, heavy quark fragmentation functions, and nucleon PDFs. Permits MC integration over phase space.



Harris, Smith 98

$F_2^{c\bar{c}}$ code: QCD-based parametrization for inclusive charm structure function

Riemersma, Smith, van Neerven 94

- Adapt simulation tools (HVQDIS, F_{2c} codes) to eN in EIC kinematics
- Assess experimental conditions for heavy quark production in eN at EIC
 - (1) Estimate charm production rates based on **generic assumptions about overall charm reconstruction efficiency at EIC.**
 - (2) Simulate **“idealized” charm reconstruction at EIC**, using PYTHIA to generate the hadronic final state, including acceptance, background, angular distributions, but assuming idealized charm reconstruction parameters
 - (3) Simulate **“realistic” charm reconstruction at EIC**, using the same setup as in (2) but including also the momentum reconstruction resolution from detector specifications.
Requires model of generic detector resolutions, to be developed as part of the project.
- Assess prospects for nuclear ratio measurements with EIC
 - Stat and sys errors, luminosity elimination, optimal choice of kinematics
 - Specifics of large $x \gtrsim 0.5$? Using beauty?

Glucos with open charm: Preliminary results



- Charm production cross section at EIC

S. Furlotov, preliminary results, 2015 APS Fall Meeting

$$0.05 < Q^2 < 1000 \text{ GeV}^2, \quad 0.5 < p_T < 10 \text{ GeV}, \quad |\eta| < 4$$

ZEUS results have different phase space, shown for comparison only

Charm cross section ~ 28 nb at $\sqrt{s} = 45$ GeV, ~ 93 nb at $\sqrt{s} = 140$ GeV


D^* cross section $\sim 30\%$ of charm cross section

- Charm production rate

~ 280 (930) charm pairs per second at luminosity $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

Rate with physics cuts will be significantly lower

Flavor separation with SIDIS



- Semi-inclusive hadron production

Standard method for charge/flavor separation of quark PDFs

HERMES, COMPASS, JLab 6 + 12 GeV

LO, NLO implementation available

- Simulation tools available: HERMES, JLab CLAS/Hall C
- Adapt to eA at EIC
- Apply to nuclear ratios at $x \sim 0.1$

Separate charge/isospin combinations with $N(\pi^+ \pm \pi^-)$

Fragmentation functions cancel in nuclear ratio. NLO effect?

Estimate final-state interactions by measuring on different nuclei and using A -dependence

- Address specific theoretical issues in nuclear ratio measurements at $x > 0.1$ and quantify physics impact
- Final-state interaction in heavy/light meson production on nuclei
 - Initial-state modifications saturate above $A \sim 12$
 - Final-state interactions continue to grow as $A^{1/3}$
 - Develop empirical method
- L/T ratio of nuclear cross sections
 - Early data indicate small nuclear modification $\Delta R/R < 0.1$ [Gousset, Pirner 96](#)
 - Revisit analysis with current data
- Impact of EIC pseudodata on PDFs
 - Bayesian reweighting: Efficient, avoids refitting

Further information

- Public Wiki at https://wiki.jlab.org/nuclear_gluons/
Simulation tools, results, materials, references
- Tools & results can be used for follow-up studies
Please contact investigators!


Supplementary material

Schedule and milestones

Months	Tasks		Milestones
1-3	Adaptation open charm MC for EIC	Furletov/Weiss	Basic MC setup for nuclear gluons with charm at EIC
4-6	Testing open charm MC for EIC Stage-1 simulations nuclear gluons Development semi-inclusive π/K MC for EIC	Furletov/Weiss/Chudakov/Stud. A Higinb./Weiss	
6-9	Stage-1 and 2 simulations nuclear gluons Testing semi-inclusive π/K MC Theoretical strategy for nuclear ratios	Furletov/Higinb./Weiss/Student A Higinb./Weiss Strikman 1 week	
9-12	Stage-2 simulations nuclear gluons Development EIC detector model for charm Theoretical update QCD description	Student A/Higinb./Weiss Hyde 4 weeks Stratmann 2 weeks Strikman 1 week	
13-15	Stage-3 simulations nuclear gluons Simulations nuclear quarks	Furletov/Higinb./Weiss Student B/Higinb./Weiss	Comprehensive assessment nuclear gluons with heavy quarks
16-18	Comprehensive assessment nuclear gluons Optimization kinematics nuclear gluons Impact global PDF fits Detailed simulations nuclear quarks Physics assessment nuclear quarks	All Strikman 1 week Stratmann 2 weeks	
19-21	Publication nuclear gluons Development EIC detector model semi-incl. π/K	All Hyde 4 weeks	
	Optimization kinematics/strategy nuclear quarks Comprehensive assessment nuclear quarks	All Strikman 1 week	
21-24	Publication nuclear quarks Overall assessment physics impact: NN interaction, non-nucleonic DOF	All	
			Comprehensive assessment nuclear quarks & gluons with EIC


MEIC detector concept

12



MEIC central detector

Ion beam from left at 50 mrad
Electron beam from right



Central & forward detectors integrated with beam optics
Ion beam from lower left
Electron beam from upper right

Information on current MEIC machine/detector design at: <https://eic.jlab.org/wiki/>