

The PRad Experiment

Maxime Levillain

September 1, 2017



The Proton Radius Puzzle

PRad Setup

Detectors Performance

Analysis

Summary

The Proton Radius Puzzle

Different Methods of Measurement

Elastic ep Scattering

New Experiment Needed

PRad Setup

Detectors Performance

Analysis

Summary

- ▶ First measurement at SLAC in 1961 through ep scattering
- ▶ 60 years of measurements, 4 possible different methods

Atomic Hydrogen Spectroscopy

Lamb shift measurements by MPQ and LKB

ep Scattering

Accelerator based experiments at Mainz, SLAC, JLab, etc


Muonic Hydrogen Spectroscopy

Lamb shift measurements by CREMA

μp Scattering

Future experiment PSI/MUSE

► Lamb shift measurements




- atomic hydrogen spectroscopy results compatible with ep scattering results
- muonic hydrogen spectroscopy results at 0.84 fm

- Elastic cross-section in the limit of the first Born approximation:

$$\frac{d\sigma}{d\Omega} = \left(\frac{d\sigma}{d\Omega} \right)_{Mott} \cdot \frac{E'}{E} \cdot \frac{1}{1+\tau} \cdot (G_E^p(Q^2) + \frac{\tau}{\epsilon} G_M^p(Q^2))$$

with:

$$Q^2 = 4EE' \sin^2 \theta / 2 \quad \tau = \frac{Q^2}{4M_p^2} \quad \epsilon = 1 / (1 + 2(1 + \tau) \tan^2 \theta / 2)$$



- Structureless proton:


$$\left(\frac{d\sigma}{d\Omega} \right)_{Mott} = \frac{\alpha^2 (1 - \beta^2 \sin^2 \theta / 2)}{4k^2 \sin^4 \theta / 2}$$

- G_E can be expressed using a Taylor expansion at low Q^2 :


$$G_E = 1 - \frac{Q^2}{6} \langle r^2 \rangle + \frac{Q^4}{120} \langle r^4 \rangle + \dots$$

which gives:

$$\langle r^2 \rangle = -6 \cdot \left. \frac{dG_E^p}{dQ^2} \right|_{Q^2=0}$$



Phys. Rev. C 93, 065207




$$r_p(e^-) = 0.8770 \pm 0.0045 \text{ fm}$$

$$r_p(\mu^-) = 0.8409 \pm 0.0004 \text{ fm}$$

- ▶ Discrepancy between muonic hydrogen spectroscopy and atomic hydrogen (spectroscopy and scattering) measurements

- ▶ Previous measurements have large systematic uncertainties and a limited coverage at small Q^2
- ▶ Requirements for PRad Experiment:
 - ▶ large Q^2 range
 - ▶ extend to very low Q^2
 - ▶ controlled systematics at sub-percent precision
- ▶ Choices:
 - ▶ Non magnetic spectrometer method
 - ▶ No target windows
 - ▶ high resolution high acceptance spectrometer
 - ▶ Normalization by Møller cross-section



Phys. Rev. C 93, 065207

- 2011 - 2012 Initial proposal
- 2012 Approved by JLab PAC39
- 2012 Funding proposal for windowless H₂ gas flow target
- 2012 - 2015 Development, construction of the target
- 2013 Funding proposals for the GEM detectors
- 2013 - 2015 Development, construction of the GEM detectors
- 2015, 2016 Experiment readiness reviews
- January/April 2016 Beam line installation
- May 2016 Beam commissioning
- May 24 - May 31 Detectors calibration
- June 4 - June 15 1.1 GeV data taking
- June 15 - June 22 2.2 GeV data taking

The Proton Radius Puzzle

PRad Setup

JLab Facility

PRad Setup

Windowless Gas Flow Target


Hybrid Calorimeter

GEM detectors

Detectors Performance


Analysis


Summary



PRad was performed in Hall B at JLab

- ▶ First experiment finished using 12 GeV accelerator (not at full beam energy)








- ▶ Electron beam or tagged photon beam at ~ 1 GeV and ~ 2 GeV
- ▶ Windowless H_2 gas flow target
- ▶ Vacuum box
- ▶ GEM detectors
- ▶ Primex HyCal

- ▶ gas target of cryogenically cooled hydrogen at 19.5 K
- ▶ beam opening: 2 mm, length: 4 cm
- ▶ cell density: $\sim 2 \cdot 10^{18}$ H atoms/cm²
- ▶ pressures:
 - ▶ cell pressure: 471 mTorr
 - ▶ chamber pressure: 2.34 mTorr
 - ▶ vacuum chamber pressure: 0.3 mTorr

Developed and build by JLab target group








- ▶ 1.7 m diameter, 2 mm aluminum vacuum window
- Limited background

Hybrid detector:

- ▶ Central part:
 - ▶ 34 x 34 matrix of PbWO_4 detectors
 - ▶ dimension of block: $2 \times 2 \times 18 \text{ cm}^3$
 - ▶ 2 x 2 blocks removed from the center for beam line to pass through
- ▶ Peripheral part:
 - ▶ 576 lead glass detectors
 - ▶ dimension of block: $4 \times 4 \times 45 \text{ cm}^3$
- ▶ Successfully used for Primex experiments



- ▶ Two large area GEM detectors: 55 cm x 123 cm
- ▶ Purpose:
 - ▶ improve spatial resolution by a factor 20 to 40 → 100 μm
 - to reduce uncertainties on θ and Q^2
- ▶ Central overlap between the 2 planes and central hole for the beam line



Developed and build by UVA

The Proton Radius Puzzle

PRad Setup

Detectors Performance

HyCal Energy Resolution

Trigger Efficiency


GEM Matching Efficiency

GEM Spatial Resolution


Analysis

Summary



- ▶ Crystal energy resolution with statistical uncertainties



- ▶ Achieved expected energy resolution:
 - ▶ 2.5% at 1 GeV for crystal part
 - ▶ 6.1% at 1 GeV for lead glass part




- ▶ Plateau from 500 MeV with an efficiency of 0.995
- ▶ Good uniformity




- ▶ GEM detection efficiency measured in both photon beam calibration (pair production) and production runs (ep and ee)
- ▶ Almost flat efficiency $> 97\%$ after removal of spacers and dead zones

X Resolution



Y Resolution



- ▶ Really good spatial resolution $\sim 74 \mu m$
- ▶ 20 to 40 times better than HyCal spatial resolution

The Proton Radius Puzzle

PRad Setup

Detectors Performance

Analysis

Stability

Yields

Cross-sections


Summary

- ▶ Calibration with tagged photon beam
 - ▶ Every calorimeter module moved into the beam
 - ▶ Allows study of resolution, linearity, trigger efficiency

- ▶ 1.1 GeV electron beam
 - ▶ 4.2 mC
 - ▶ 604 M events with target
 - ▶ 53 M events with “empty target”
 - ▶ 25 M events with ^{12}C target for calibration

- ▶ 2.2 GeV electron beam
 - ▶ 14.3 mC
 - ▶ 756 M events with target
 - ▶ 38 M events with “empty target”
 - ▶ 10.5 M events with ^{12}C target for calibration



- ▶ Control of target properties (pressure, temperature, position) via EPICS




→ Less than 3% deviation

Weizhi Xiong

- Phase space after background subtraction





- Separation of ep and Møller phase space (for $\theta > 0.85$ deg for 1 GeV)



- ▶ 1.1 GeV data set: $Q^2 \in [2 \cdot 10^{-4}, 1.3 \cdot 10^{-2}] \text{ GeV}^2$
- ▶ 2.2 GeV data set: $Q^2 \in [8 \cdot 10^{-4}, 6 \cdot 10^{-2}] \text{ GeV}^2$

- ▶ Stability of ratio ep/ee after background subtraction for different beam intensity



- ▶ Good stability for the 2GeV period


- ▶ Normalization of ep cross-section by Møller cross-section:

$$\left(\frac{d\sigma}{d\Omega}\right)_{ep} = \frac{N_{exp}(ep \rightarrow ep \text{ in } \theta_i \pm \Delta\theta)}{N_{exp}(ee \rightarrow ee)} \cdot \frac{\epsilon_{geom}^{ee}}{\epsilon_{geom}^{ep}} \cdot \frac{\epsilon_{det}^{ee}}{\epsilon_{det}^{ep}} \cdot \left(\frac{d\sigma}{d\Omega}\right)_{ee}$$


- ▶ Several event generators have been developed for ep and Møller scattering taking into account complete calculations of radiative corrections beyond ultra relativistic approximations
 - ▶ A. V. Gramolin et al., J. Phys. G Nucl. Part. Phys. 41(2014)115001
 - ▶ I. Akushevich et al., Eur. Phys. J. A 51(2015)1
- ▶ Geant4 is used to take into account all external radiative effects

$$\sigma_{ep}^{Born} = \left(\frac{\sigma_{ep}}{\sigma_{ee}}\right)^{exp} / \left(\frac{\sigma_{ep}}{\sigma_{ee}}\right)^{sim} \cdot \sigma_{ee}^{Born}$$

- ▶ Preliminary ep cross-section for the 2.2 GeV data set
- ▶ Statistical uncertainties at $\sim 0.2\%$ per point
- ▶ Conservative point-to-point systematic uncertainties at $\sim 2\%$



- ▶ Preliminary ep cross-section for the 2.2 GeV data set
- ▶ Statistical uncertainties at $\sim 0.2\%$ per point
- ▶ Conservative point-to-point systematic uncertainties at $\sim 2\%$



- ▶ The PRad experiment was uniquely designed to address the *Proton Radius Puzzle*
- ▶ The experiment was successfully performed in May-June 2016
- ▶ Wide range of Q^2 without normalization on more than two orders of magnitude ($2 \cdot 10^{-4} \text{ GeV}^2$ to $6 \cdot 10^{-2} \text{ GeV}^2$)
- ▶ Lowest Q^2 data set of ep elastic scattering ($2 \cdot 10^{-4} \text{ GeV}^2$)
- ▶ First preliminary extraction of the proton radius expected at the end of October

Thanks to JLab, Hall B, Accelerator Division and Target Group

PRad is supported in part by NSF MRI award PHY-1229153, as well as DOE awards for GEM; my research work is supported by NSF awards: PHY-1506388 and PHY-0855543