

Status of PRad Experiment

Chao Gu

Duke University


For PRad Collaboration

Outline

- The Proton Charge Radius
- Experiment Setup
- Analysis Status and Preliminary Results

The Proton Charge Radius Puzzle

- Proton radius is one of the most fundamental quantities in physics:
 - Critically important for atomic physics in precision spectroscopy of atom
 - Precision test of nuclear/particle models
 - Connects atomic and subatomic physics




R. Pohl et al., Nature 466, 213 (2010)

A. Antognini et al., Science 339, 417 (2013)

The Proton Charge Radius Puzzle

- 4 different methods to measure the proton charge radius:
 - Hydrogen spectroscopy (ordinary hydrogen, muonic hydrogen)
 - Lepton-proton elastic scattering (ep, μp)
- The proton charge radius puzzle:
 - $\sim 8\sigma$ discrepancy between the new muonic-hydrogen spectroscopy measurements and all previous results



R. Pohl et al., Nature 466, 213 (2010)



A. Antognini et al., Science 339, 417 (2013)

ep Scattering

- Previous measurements have large systematic uncertainties and a limited coverage at small Q^2
- Requirements for PRad Experiment:
 - large Q^2 range
 - extend to very low Q^2
 - controlled systematics at sub-percent precision

- Extraction of $\langle r^2 \rangle = -6 \left. \frac{dG_E^p(Q^2)}{dQ^2} \right|_{Q^2=0}$

$$\frac{d\sigma}{d\Omega} = \left(\frac{d\sigma}{d\Omega} \right)_{\text{Mott}} \left(\frac{E'}{E} \right) \frac{1}{1 + \tau} \left(G_E^2(Q^2) + \frac{\tau}{\varepsilon} G_M^2(Q^2) \right)$$




PRad Timeline

- 2011 - 2012 Initial proposal
- 2012 Approved by JLab PAC39
- 2012 Funding proposal for windowless H₂ gas flow target
- 2012 - 2015 Development, construction of the target
- 2013 Funding proposals for the GEM detectors
- 2013 - 2015 Development, construction of the GEM detectors
- 2015, 2016 Experiment readiness reviews
- Jan - Apr 2016 Beam line installation
- May 2016 Beam commissioning
- May 24 - 31 2016 Detectors calibration
- Jun 4 - 22 2016 Data taking


PRad Setup

- Electron beam at 1.1 GeV and 2.2 GeV
- Windowless H₂ gas flow target
- Vacuum box
- GEM detectors
- PrimEx HyCal




Windowless H₂ Gas Flow Target

- A windowless gas target of cryogenically cooled hydrogen:
 - 4 cm long copper target cell
 - 7.5 μm kapton windows with 2 mm beam orifices
 - H₂ gas cooled at 19.5 K
 - Target density: $\sim 2 \times 10^{18}$ H atoms/cm²
 - Four-axis motion system to position the target cell with 10 μm accuracy
- Pressures:
 - Cell pressure: 471 mTorr
 - Chamber pressure: 2.34 mTorr
 - Vacuum chamber pressure: 0.3 mTorr
- Two additional solid target foils: 1 μm ¹²C and Al





Vacuum Box

- 1.7 m diameter, 1.6 mm aluminum vacuum window




Electromagnetic Calorimeter (PrimEx HyCal)

- Combination of PbWO_4 and Pb-glass detectors ($118 \times 118 \text{ cm}^2$)
 - 34×34 matrix of $2.05 \times 2.05 \times 18 \text{ cm}^3$ PbWO_4 shower detectors
 - 576 Pb-glass shower detectors ($3.82 \times 3.82 \times 45 \text{ cm}^3$)
 - 2×2 PbWO_4 modules removed in the middle for beam passage
- 5.8 m from the target
- Successfully used for PrimEx experiments



GEM Detectors

- Two large area GEM detectors (55 cm x 123 cm)
 - Central overlapped between two planes with a hole for the beam passage
- Purpose:
 - Factor of >20 improvements in coordinate resolutions
 - Similar improvements in Q^2 resolution (important)
 - Unbiased coordinate reconstruction
 - Increase Q^2 coverage by including HyCal Pb-glass part
- Designed and built at University of Virginia (UVa)






Experiment Data Collected

- With 1.1 GeV beam:
 - Collected 4.2 mC
 - 604 M events with H₂ target
 - 53 M events with “empty” target
 - 25 M events with ¹²C target for calibration
- With 2.2 GeV beam:
 - Collected 14.3 mC
 - 756 M events with H₂ target
 - 38 M events with “empty” target
 - 10.5 M events with ¹²C target for calibration



GEM Resolution

- Extraction of GEM spatial resolution using GEM central overlapping region
- Good spatial resolution achieved




GEM Efficiency

- GEM efficiency calibrated using physics runs
 - GEM spacer introduces deficient area
 - Evenly distributed efficiency after spacer correction
- Stable GEM efficiency over time
 - Average efficiency fluctuation: $\sim 0.5\%$ level



HyCal Calibration


- Gains controlled by Light Monitoring System (LMS)
- Two different calibrations:
 - Before data taking: Scan with 250–1050 MeV tagged photon beam moved in front of each module to study of resolution, efficiency and non linearity
 - During data taking: With Moller and ep events
- Achieved expected energy resolution:
 - 2.5% at 1 GeV for PbWO₄ part
 - 6.1% at 1 GeV for Pb-glass part
- Plot shows the energy resolution for PbWO₄ part with statistical uncertainties and systematic coming from non-uniformity




Preliminary Results

- The plot shows the extracted differential cross section vs scattered angle, for 2.2 GeV incident beam energy in 0.7 - 3.5 deg range (Very Preliminary)
- Statistical errors are ~0.2% per point
- Systematic errors at this stage are estimated to be on 2%(?) level (shown with a dashed area).

ep elastic scattering cross section



ep elastic scattering cross section



Conclusion

- The PRad experiment was uniquely designed to address the “Proton Radius Puzzle”
- Experiment had been successfully performed in May - June, 2016
- About half of the 2.2 GeV beam energy data have been analyzed so far:
 - Very preliminary differential cross sections for the elastic ep scattering have been extracted for the forward angle range from 0.70 to 3.00 deg
- Preliminary fit to extract the proton radius from the 2.2 GeV data set is expected to be done by October this year

Thanks