







OHIO  
UNIVERSITY

# Photoproduction of scalar mesons with g12 data



---

# gl2 : Event selection



Threshold photon energy for production of  $f_0(1500)$  is 2.7  $\text{GeV}$

# $K_s^0 K_s^0$ Invariant mass spectrum




- Timing cuts for pion identification
- Standard eloss corrections
- Cut on Missing mass (proton)
- $E_T > 2.7$  GeV
- $M(K_s)$  is within 16 MeV of the PDG value

The  $K_s^0 K_s^0$  invariant mass spectrum has a peak around 1280 MeV and another around 1500 MeV.


The 1280 peak, which may be the  $f_2(1270)/a_2(1320)$  has a much narrower width than the PDG average value. Peak at 1500 either  $f_0(1500)$  or  $f'_2(1525)$

# $K^0_s K^0_s$ Invariant mass spectrum




# Dalitz Plots

Dalitz plot of  $K_S K_S$  vs  $K_S p$ ,  $|t| < 1$



Dalitz plot of  $K_S K_S$  vs  $K_S p$ ,  $|t| > 1$



$$\langle Y_{00} \rangle = N \left[ |S|^2 + |P_-|^2 + |P_0|^2 + |P_+|^2 + |D_-|^2 + |D_0|^2 + |D_+|^2 \right]$$

$$\langle Y_{10} \rangle = N \left[ SP_0^* + P_0 S^* + \sqrt{\frac{3}{5}} (P_- D_-^* + P_- S^* + P_+ D_+^* + D_+ P_+^*) + \sqrt{\frac{4}{5}} (P_0 D_0^* + D_0 P_0^*) \right]$$


$$\begin{aligned} \langle Y_{11} \rangle = N & \left[ \sqrt{\frac{1}{2}} (-P_- S^* - S P_-^* + P_+ S^* + S P_+^*) + \sqrt{\frac{1}{20}} (P_- D_0^* + D_0 P_-^* - P_+ D_0^* - D_0 P_+^*) \right. \\ & \left. + \sqrt{\frac{3}{20}} (-P_0 D_-^* - D_- P_0^* + P_0 D_+^* + D_+ P_0^*) \right] \end{aligned}$$












$$\begin{aligned} \langle Y_{20} \rangle = N & \left[ SD_0^* + D_0 S^* + \sqrt{\frac{1}{5}} (2|P_0|^2 - |P_-|^2 - |P_+|^2 + |F_-|^2 + |F_+|^2) \right. \\ & \left. + \sqrt{\frac{5}{49}} (|D_-|^2 + |D_+|^2) + \sqrt{\frac{20}{49}} |D_0|^2 \right] \end{aligned}$$


$$\begin{aligned} \langle Y_{21} \rangle = N & \left[ \frac{1}{2} (SD_+^* + D_+ S^* - SD_-^* - D_- S^*) + \sqrt{\frac{3}{20}} (P_0 P_+^* + P_+ P_0^* - P_- P_0^* - P_0 P_-^*) \right. \\ & \left. + \sqrt{\frac{5}{196}} (D_0 D_+^* + D_+ D_0^* - D_0 D_-^* - D_- D_0^*) \right] \end{aligned}$$


$$\langle Y_{22} \rangle = N \left[ \sqrt{\frac{3}{10}} (P_- P_+^* + P_+ P_-^*) + \sqrt{\frac{3}{196}} (-D_- D_+^* - D_+ D_-^*) \right]$$

# Moments Analysis










## Ks angles for pure D0, in Gottfried Jackson frame of resonance X


**cosθ, pureD fullAcc, 1500-1550MeV**



**ctgj**

Entries 438292  
Mean 0.0004803  
RMS 0.7231







**φ, pureD fullAcc, 1500-1550MeV**



**phigj**

Entries 438292  
Mean -0.000112  
RMS 0.5771

Pure D0 , full acceptance, no cuts on t



$$\langle Y_{00} \rangle = N \left[ |S|^2 + |P_-|^2 + |P_0|^2 + |P_+|^2 + |D_-|^2 + |D_0|^2 + |D_+|^2 \right]$$

$$\langle Y_{10} \rangle = N \left[ SP_0^* + P_0 S^* + \sqrt{\frac{3}{5}} (P_- D_-^* + P_- S^* + P_+ D_+^* + D_+ P_+^*) + \sqrt{\frac{4}{5}} (P_0 D_0^* + D_0 P_0^*) \right]$$

$$\begin{aligned} \langle Y_{11} \rangle = N & \left[ \sqrt{\frac{1}{2}} (-P_- S^* - S P_-^* + P_+ S^* + S P_+^*) + \sqrt{\frac{1}{20}} (P_- D_0^* + D_0 P_-^* - P_+ D_0^* - D_0 P_+^*) \right. \\ & \left. + \sqrt{\frac{3}{20}} (-P_0 D_-^* - D_- P_0^* + P_0 D_+^* + D_+ P_0^*) \right] \end{aligned}$$

$$\begin{aligned} \langle Y_{20} \rangle = N & \left[ SD_0^* + D_0 S^* + \sqrt{\frac{1}{5}} (2|P_0|^2 - |P_-|^2 - |P_+|^2 + |F_-|^2 + |F_+|^2) \right. \\ & \left. + \sqrt{\frac{5}{49}} (|D_-|^2 + |D_+|^2) + \sqrt{\frac{20}{49}} |D_0|^2 \right] \end{aligned}$$

$$\begin{aligned} \langle Y_{21} \rangle = N & \left[ \frac{1}{2} (SD_+^* + D_+ S^* - SD_-^* - D_- S^*) + \sqrt{\frac{3}{20}} (P_0 P_+^* + P_+ P_0^* - P_- P_0^* - P_0 P_-^*) \right. \\ & \left. + \sqrt{\frac{5}{196}} (D_0 D_+^* + D_+ D_0^* - D_0 D_-^* - D_- D_0^*) \right] \end{aligned}$$

$$\langle Y_{22} \rangle = N \left[ \sqrt{\frac{3}{10}} (P_- P_+^* + P_+ P_-^*) + \sqrt{\frac{3}{196}} (-D_- D_+^* - D_+ D_-^*) \right]$$

# Moments Analysis

# Pure D0 , full acceptance

